

BEST AVAILABLE COPY

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁵ : C12N 15/00, 7/01, 5/10 C12P 21/02</p>	<p>A1</p>	<p>(11) International Publication Number: WO 90/05780 (43) International Publication Date: 31 May 1990 (31.05.90)</p>
<p>(21) International Application Number: PCT/US89/05285 (22) International Filing Date: 20 November 1989 (20.11.89) (30) Priority data: 273,373 18 November 1988 (18.11.88) US (71) Applicant: STATE OF OREGON, acting by and through THE OREGON STATE BOARD OF HIGHER EDUCATION, acting for and on behalf of THE OREGON HEALTH SCIENCES UNIVERSITY [US/US]; 3181 S.W. Sam Jackson Park Road, L-335, Portland, OR 97201 (US). (72) Inventors: CIVELLI, Olivier ; 3012 N.E. 24th Avenue, Portland, OR 97212 (US). BUNZOW, James, R. ; 9904 S.W. 56th Avenue, Portland, OR 97219 (US). GRANDY, David, K. ; 3730 S.W. Patton Road, Portland, OR 97221 (US). MACHIDA, Curtis, A. ; 12720 N.W. Lovejoy, Portland, OR 97229 (US).</p>		<p>(74) Agents: ZELANO, Anthony, J. et al.; Millen, White & Zelano, Arlington Courthouse Plaza I, Suite 1201, 2200 Clarendon Boulevard, Arlington, VA 22201 (US). (81) Designated States: AT (European patent), AU, BE (European patent), CH (European patent), DE (European patent), ES (European patent), FR (European patent), GB (European patent), IT (European patent), JP, LU (European patent), NL (European patent), SE (European patent). Published With international search report.</p>
<p>(54) Title: DOPAMINE RECEPTORS AND GENES</p> <p>(57) Abstract</p> <p>A mammalian D₂ dopamine receptor gene has been cloned. Thus, DNA sequences encoding all or a part of the dopamine receptor are provided, as well as the corresponding polypeptide sequences and methods for producing the same synthetically and via expression of a corresponding sequence from a host transformed with a suitable vector carrying the corresponding DNA sequence. The various structural information provided by this invention enables the preparation of labeled or unlabeled immunospecific species, particularly antibodies, as well as nucleic acid probes labeled in conventional fashion. Pharmaceutical compositions and methods of using various products of this invention are also provided.</p>		

13 A

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MR	Mauritania
BE	Belgium	GA	Gabon	MW	Malawi
BF	Burkina Faso	GB	United Kingdom	NL	Netherlands
BG	Bulgaria	HU	Hungary	NO	Norway
BJ	Benin	IT	Italy	RO	Romania
BR	Brazil	JP	Japan	SD	Sudan
CA	Canada	KP	Democratic People's Republic of Korea	SE	Sweden
CF	Central African Republic	KR	Republic of Korea	SN	Senegal
CG	Congo	LI	Liechtenstein	SU	Soviet Union
CH	Switzerland	LK	Sri Lanka	TD	Chad
CM	Cameroon	LU	Luxembourg	TG	Togo
DE	Germany, Federal Republic of	MC	Monaco	US	United States of America
DK	Denmark				

DOPAMINE RECEPTORS AND GENES**Background of the Invention**

This invention relates to dopamine receptors from mammalian species and the corresponding genes. In particular, it relates to the isolation, sequencing and/or cloning of D₂ dopamine receptor genes, the synthesis of D₂ dopamine receptors by transformed cells, and the manufacture and use of a variety of novel products enabled by the identification and cloning of DNA encoding dopamine receptors.

Dopamine receptors in general have been implicated in a large number of neurological and other disorders, including, for example, movement disorders, schizophrenia, drug addiction, Parkinson's disease, Tourette syndrome, Tardive Dyskinesia, and many others. As a result, the dopamine receptor has been the subject of numerous pharmacological and biochemical studies.

In general, dopamine receptors can be classified into D₁ and D₂ subtypes based on pharmacological and biochemical characteristics (1,2). The D₂ dopamine receptor has been implicated in the pathophysiology and treatment of the mentioned disorders. In addition, it is known that the D₂ dopamine receptor interacts with guanine nucleotide binding proteins to modulate second messenger systems (6,7).

Despite the heavy emphasis placed on elucidation of the existence and properties of the dopamine receptor and its gene, identification, isolation and cloning have been inaccessible heretofore. This is true despite the knowledge that other members of the family of receptors that are coupled to G proteins share a significant

similarity in primary amino acid sequence and exhibit an archetypical topology predicted to consist of seven putative transmembrane domains (8). Regarding the serotonin receptor, e.g., see Julius et al., Science, 5 Vol. 241, 558 (1988).

Summary of the Invention

This invention has successfully identified, isolated and cloned mammalian, including human, D₂ dopamine receptor gene sequences and produced the encoded dopamine receptor. The corresponding polypeptide has been synthesized. Sequences of both the gene and the polypeptide have been determined. The invention also provides a variety of new and useful nucleic acid, cell line, vector, and polypeptide and medicinal products, inter alia, as well as methods of using these.

Thus, this invention relates to an isolated DNA sequence, an identified portion of which is a structural gene which encodes a polypeptide having the biological activity of a mammalian D₂ dopamine receptor. In particular, it relates to an isolated DNA sequence which will hybridize to a DNA sequence encoding a mammalian D₂ dopamine receptor. It also relates to fragments, variants and mutants of such sequences, particularly those which also encode a polypeptide having biological activity of a mammalian dopamine receptor, most particularly a mammalian D₂ dopamine receptor. In a preferred aspect, the dopamine receptor is human. In another preferred aspect, the sequence is that of rat D₂ dopamine receptor as shown in Figure 1. Of course, the nucleic acids of this invention also include complementary strands of the foregoing, as well as sequences differing therefrom by codon degeneracy and sequences which hybridize with the aforementioned sequences.

In other preferred aspects, this invention includes nucleic acid sequences and fragments useful as

oligonucleotide probes, preferably labelled with a detectable moiety such as a radioactive or biotin label. For example, such probes can hybridize with DNA encoding a polypeptide having the biological activity of a D₂ dopamine receptor or with DNA associated therewith, e.g., DNA providing control of a D₂ dopamine receptor gene or introns thereof, etc. DNA of this invention can also be part of a vector.

The invention also involves cells transformed with vectors of this invention as well as methods of culturing these cells to manufacture polypeptides, e.g., having the biological activity of a D₂ dopamine receptor. Preferably, the cells are of mammalian origin when used in such methods.

The invention also relates to polypeptides encoded by the foregoing nucleic acid sequences, especially to isolated mammalian dopamine receptors, preferably of human origin. The invention further relates to polypeptides which are mutants or variants of such receptors, preferably those wherein one or more amino acids are substituted for, inserted into and/or deleted from the receptor, especially those mutants which retain the biological activity of a dopamine receptor. This invention also relates to antibodies, preferably labelled, and most preferably monoclonal, capable of binding a dopamine receptor amino acid sequence, preferably wherein the latter is human, or a fragment of such an antibody.

The invention further relates to compositions comprising one of the various products mentioned above and, typically, a pharmaceutically acceptable carrier as well as to methods of employing these products to achieve a wide variety of utilitarian results.

Brief Description of the Drawings

Various other objects, features and attendant advantages of the present invention will be more fully appreciated as the same becomes better understood when considered in conjunction with the accompanying drawings, and wherein:

Figure 1 shows the nucleotide sequence for the RGB-2 cDNA and deduced amino acid sequence of the rat D₂ dopamine receptor. The nucleotide sequence is numbered beginning with the first methionine of the long open reading frame. Nucleotide numbering appears beneath the nucleotide sequence at the right-hand end of each line. The deduced amino acid sequence is shown above the nucleotide sequence. The double underline denotes the small open reading frame in the 5' untranslated region. Postulated N-linked glycosylation sites are indicated by asterisks. Putative protein kinase A phosphorylation sites have a line above them. The intron splice junction is designated by an arrow. The poly A adenylation site is underlined.

Figure 2 shows the alignment of the amino acid sequences of the rat D₂ dopamine receptor, the hamster β₂-adrenergic receptor, the human α₂-adrenergic receptor, the human G-21 (serotonin 1a) receptor, the porcine muscarinic M receptor and the bovine substance K receptor. Amino acids enclosed in solid lines and shaded represent residues that are identical in at least two other sequences when compared to the D₂ dopamine receptor. The putative transmembrane domains are indicated by brackets labeled by Roman numerals. The numbers of residues in the variable third cytoplasmic loop and at the C-terminus are in parentheses.

Figure 3 shows a Northern blot analysis of RGB-2 transcripts in rat brain and pituitary. Each lane contained 20 μg of total RNA. Numbers on the right indicate molecular weight as determined from RNA size markers (BRL). A: olfactory bulb; B: hippocampus; C: cere-

bellum; D: posterior cortex; E: anterior cortex; F: thalamus; G: hypothalamus; H: medulla; I: amygdala; J: mesencephalon; K: septum; L: posterior basal ganglia; M: anterior basal ganglia; N: neurointermediate lobe of pituitary; O: anterior lobe of the pituitary.

Figure 4 illustrates the binding of ^3H -spiperone to membranes from L-RGB2Zem-1 cells.

a) 1: Saturation isotherms of the specific binding of ^3H -spiperone to membranes prepared from L-RGB2Zem-1 cells and rat striatum. Results are shown from one of four independent experiments.

2: Scatchard transformation of the data.

b) Competitions curves using L-RGB2Zem-1 membranes. Representative curves are shown for inhibition of specifically bound ^3H -spiperone by drugs in membranes from L-RGB2Zem-1 cells. Each drug was tested 3 times.

c) Table of K_i values for L-RGB2Zem-1 and rat striatum. Results are geometric means of 3 experiments in which 0.5 nM ^3H -spiperone was inhibited by various concentrations of unlabeled drug. For some drugs, inhibition curves in rat striatal tissue were fit best by assuming the presence of two classes of binding sites. The proportions of binding sites with high or low affinity for inhibitor are shown in parentheses. K_i values for the class of binding sites representing 10-25% of specific binding were calculated by assuming that the radioligand was binding to serotonin receptors with a K_d value of 1 nM.

Figure 5 shows (a) a hydrophobicity plot of the amino acid sequence shown in Figure 1; and (b) a hydrophobicity plot of the amino acid sequence of the β_2 -adrenergic receptor. The transmembrane regions are marked by the Roman numerals.

Figure 6 shows a calculated restriction map of a 2477 base EcoRI fragment of the nucleic acid sequence shown in Figure 1.

Figure 7 shows a partial sequence of a human D₂ dopamine receptor, the middle amino acid sequence shown being the correct one.

Figure 8 shows a saturation analysis of specific [3H]spiroperidol binding to LZR1 and LZR2 cells. Results shown are from representative experiments in which LZR1 and LZR2 cells were grown in control growth medium or medium to which zinc sulfate had been added, as indicated. Data are plotted as specifically bound radioligand divided by the corrected free concentration of radioligand (total added minus total bound), versus specifically bound radioligand. For zinc treatments, 100 µM zinc sulfate was added to growth medium for 16 hours. Both control and zinc-treated cells were then washed and grown in control medium for one day before harvesting. In the experiments shown, B_{max} values determined by nonlinear regressions analysis were 876 and 914 fmol/mg of protein for control LZR1 cells or LZR1 cells treated with zinc, respectively. B_{max} values for control or zinc-treated LZR2 cells were 385 and 593 fmol/gm of protein, respectively.

Figure 9 shows the inhibition of radioligand binding by agonists. Results are plotted as specific binding, expressed as a percentage of specific binding in the absence of competing drug, versus the log of the concentration of competing drug. Membranes were prepared from LZR1 cells as described in the text.

A. Curves from a single experiment are shown for inhibition of the binding of [3H]spiroperidol by agonists. Each drug was tested twice. In this experiment, the free concentration of [3H]spiroperidol was 230 pM, and the K_b value for [3H]spiroperidol was 60 pM. K_i values and Hill coefficients in this experiment were 5 nM and 1.05 for bromocriptine, respectively, 790 nM and .89 for (-)3-PPP, 8 µM and 1.0 for quinpirole, 31 µM and 1.05 for (+)3-PPP, and 0.3 mM and 0.72 for LY181990.

B. Results are shown from one of four independent experiments in which the effect of GTP and NaCl on the inhibition of [³H]spiroperidol binding by DA was determined. Concentrations of [³H]spiroperidol ranged from 323 to 498 pM. In this experiment, the concentration of radioligand was 323 pM. Open circles represent inhibition by DA in the presence of 0.1 mM GTP and 120 mM NaCl, whereas closed circles represent inhibition in the absence of added GTP and NaCl. IC₅₀ values and Hill coefficients in this experiment were 29 μ M and 0.65, respectively, in the absence and 115 μ M and 1.03 in the presence of GTP and NaCl.

Figure 10 shows the inhibition of adenylate cyclase activity in LZR1 cells. Agonists were tested for inhibition of adenylate cyclase activity in membranes prepared from LZR1 cells. Approximately 50 to 100 μ g of protein was used in each assay. Results are shown as [³²P]cAMP/mg of protein/min., expressed as a percentage of total activity in the presence of 10 μ M forskolin. Representative dose-response curves are shown for six drugs, each tested at least three times. Data are plotted as enzyme activity versus the log of the concentration of drug. No curve is plotted for the data for (-)-3-PPP, since no inhibition was observed. In the experiments shown in this figure, basal and forskolin-stimulated activity ranged from approximately 0.8 to 1.5 and 8.5 to 15.8 pmol/mg of protein/min., respectively.

Figure 11 shows the blockade of DA-sensitive adenylate cyclase. Results shown are means of three experiments \pm SE, plotted as the percentage of total activity in the presence of 10 μ M forskolin. Forskolin (FSK) was present in all the experiments shown, together with 10 μ M dopamine (DA) or DA and 10 μ M (+)-butaclamol (BUT) as indicated. Some cells were treated with pertussins toxin (PT) before harvesting for determination of enzyme activity. Basal activity in control and PT-treated cells was 1.2 ± 0.07 and 1.6 ± 0.15 pmol/mg of

protein/min., respectively. Total forskolin-stimulated activity in control cells (FSK) was 11.9 ± 1.0 pmol/mg of protein/min. * $p < 0.05$ compared to FSK in control cells, as determined by a t test for paired means.

5 Figure 12 shows the reversal of dopamine inhibition by pertussis toxin pretreatment. Data presented for membrane adenylate cyclase activity represent means (x) with standard error (S.E.) and % inhibition (% Inh.) below. % Inhibition was calculated from the equation
10 $100 \times [1 - (S - B / I - B_i)]$ where B, S, and I are values of basal activity, activity in the presence of stimulator (S) or inhibitor (I), respectively and B_i is basal activity in the presence of inhibitor. Results were obtained from parallel assays in controls and cells pretreated (16h, 50
15 ng/ml) with pertussis toxin (indicated as +P.T.).

(A) Adenylate Cyclase. Membranes for cyclase assay were exposed acutely to 10 μ M forskolin (FSK) or 100 μ M dopamine (DA), and adenylate cyclase activity expressed as pmol/mg protein/min.

20 (B) Intracellular cAMP. Cells were treated acutely with VIP (200 nM) and dopamine (1 μ M) and cAMP accumulation (expressed as pmol/dish) was measured in cell extracts.

(C) Extracellular cAMP. Media samples from the
25 same dishes of cells were assayed for cAMP accumulation expressed as pmol/dish.

Figure 13 shows the expression of specific dopamine- D_2 receptor mRNA and specific binding in GH_4ZR_7 transfectant cells.

30 (A) Northern blot analysis of GH_4C_1 cell total RNA (20 μ g/lane). Y-axis indicate the migration of RNA molecular weight standards (kb).

(B) (1) Specific binding of 3H -spiperone to membranes prepared from GH_4ZR_7 cells was characterized by
35 saturation analysis (see "Experimental Procedures", Example 2). Data from one of four independent experiments are plotted as specifically bound radioligand

(ordinate) versus corrected free radioligand concentration (total added minus total bound). Calculated K_D and B_{max} values for this experiment were 60 pM and 1165 fmol/mg protein.

5 (B) (2) Transformation of the data by the method of Scatchard which are plotted as specific bound/free (Y-axis) vs. specific bound concentrations of 3H -spiperone (X-axis).

10 (C) Displacement of specific 3H -spiperone binding by dopamine: effect of GTP/NaCl. GH_4ZR_7 cell membranes were incubated with 3H -spiperone (0.47 nM) and indicated concentrations of dopamine (X-axis) in the absence (o) or presence (•) of 100 μM GTP/120 mM NaCl. Results are shown for one of four experiments.

15 Calculated IC_{50} and Hill coefficient values for dopamine in the experiment shown were 16 μM and 0.61 in the absence of GTP/NaCl, and 56 μM and 0.85 in the presence of GTP/NaCl.

20 Figure 14 shows the inhibition of cAMP accumulation and PRL release by dopamine in GH_4ZR_7 cells. Incubations were performed in triplicate as described in "Experimental Procedures", Example 2.

25 (A) Inhibition of extracellular cAMP accumulation by dopamine. Parallel dishes of GH_4C_1 and GH_4ZR_7 cells were incubated with concentrations of VIP, dopamine (D), and (-)-sulpiride (-S) of 250 nM, 10 μM and 5 μM , respectively. Untreated controls are denoted as "C". Media were collected and assayed for cAMP (ordinate) expressed as pmol/dish.

30 (B) Inhibition of intracellular cAMP accumulation by dopamine in GH_4ZR_7 cells. Cell extracts were assayed for cAMP, expressed on the ordinate. Drug concentrations were as in (A), except (+)-sulpiride (+S), 5 μM .

35 (C) Inhibition of stimulated PRL release by dopamine in GH_4ZR_7 cells. Media samples were assayed for PRL (ordinate) after the indicated treatments. The

concentrations of VIP, TRH, dopamine (D), and (-)-sulpiride (-S) were 200 nM, 200 nM, 100 nM, and 2 μ M, respectively.

5 Figure 15 shows dose-response relations for dopamine inhibition of basal and VIP-enhanced cAMP accumulation in GH₄ZR₇ cells.

(A) Basal intra- (•) and extracellular (o) cAMP accumulation in the presence of indicated concentrations of dopamine. Basal cAMP levels in the absence of
10 dopamine were 22 ± 6 pmol/dish (intracellular) and 12.4 ± 0.6 pmol/dish (extracellular). EC₅₀ values for dopamine actions were 4.9 nM (intracellular) and 8.5 nM (extracellular).

(B) VIP-enhanced intra- (•) and extracellular (o) cAMP accumulation in the presence of indicated
15 dopamine concentrations. VIP (250 nM)-enhanced levels of intra- and extracellular cAMP (in the absence of dopamine) were 145 ± 1.2 pmol/dish and 146 ± 2.8 pmol/dish, and basal cAMP levels were 35 ± 1.6 pmol/dish and 15 ± 0.2
20 pmol/dish, respectively. EC₅₀ values for dopamine inhibition were 5.5 nM (intracellular) and 5.8 nM (extracellular).

Figure 16 shows the specific blockade of dopamine-induced inhibition of VIP-enhanced cAMP accumulation.
25 GH₄ZR₇ cells were incubated in FAT medium (see "Methods", Example 2) in the presence of 250 nM VIP and 100 nM dopamine, and indicated concentrations of various dopamine antagonists, and extracellular levels of cAMP measured. Data are plotted as percent of maximal cAMP
30 levels versus the logarithm of indicated concentrations of dopamine antagonists. The standard error of triplicate determinations was less than 8%. Basal and VIP-enhanced cAMP levels (in the absence of dopamine and antagonists) were 25.8 ± 1.6 pmol/dish and 214 ± 14
35 pmol/dish. IC₅₀ and estimated K_i values for antagonism of dopamine actions were: spiperone, 0.56 nM and 30 pM; (+)-butaclamol, 4.5 nM and 0.2 nM; (-)-sulpiride 38 nM

and 1.8 nM; SCH23390, >1 μ M; (-)-butaclamol, >10 μ M. Estimated K_i were calculated from the equation $K_i = IC_{50}/(1 + ([DA]/EC_{50}))$, where [DA] is 100 nM and the EC_{50} for dopamine is 6 nM. Spiperone, (+)-butaclamol and (-)-sulpiride alone did not alter basal cAMP levels.

Figure 17 shows the inhibition of adenylyl cyclase by dopamine- D_2 agonists. Inhibition of adenylyl cyclase activity was assessed in the presence of 10 μ M forskolin. Data are plotted as the mean of triplicate assays, with enzyme activity expressed as a percentage of total activity versus the logarithm of drug concentration. Average basal adenylyl cyclase activity was 4.6 ± 0.2 pmol/mg protein/min and total forskolin-stimulated activity was 63.8 ± 0.2 pmol/mg protein/min. EC_{50} values and maximal inhibition for the experiments shown were 79 nM and 57%, respectively, for dopamine, 200 nM and 49% for quinpirole, 5 nM and 23% for bromocryptine, and 600 μ M and 40% for (+)-3-PPP.

Figure 18 shows the nucleotide sequence of the human pituitary dopamine D_2 receptor cDNA. The deduced amino acid sequence is indicated above the human cDNA. Below is the nucleotide sequence of the cloned rat cDNA (see Reference 9, Example 3) and the amino acids which differ between the two clones. Boxed regions, numbered I-VII, represent the putative transmembrane domains. Triangles indicate the exon/intron splice junctions; a period is one missing base pair; asterisks identify potential N-linked glycosylation sites and targets of protein kinase A phosphorylation are underlined. The polyadenylation signal is double underlined.

Figure 19 shows competition curves of [3 H]-domperidone binding to L-hPitD $_2$ Zem membranes. The radioligand was used at a concentration of 1 nM, and specific binding was defined using 1 μ M (+) butaclamol. Data are shown for one of three experiments.

Figure 20 shows Southern blot of human genomic DNA. The genomic BamHI 1.6-kb fragment containing exon 7 was

prepared from λ HD2G1 and used as probe (specific activity, 2×10^8 cpm). DNA was digested with BamHI (lane 1), BglIII (lane 2), BamHI/BglIII (lane 3), and HindIII (lane 4).

5 Figure 21 is a schematic representation of the human dopamine D_2 receptor gene and pituitary cDNA.

(a) Restriction map of the two overlapping genomic phase, λ HD2G1 and λ HD2G2; A, ApaI; B, BamHI; Bg, BglIII; H, HindIII.

10 (b) Diagram of the human gene locus DRD2. Exons, indicated by the boxes, are numbered. The solid boxes indicate regions of coding sequence and open boxes, non-translated sequence. The genomic sequencing strategy is expanded above the gene. These regions are not drawn to scale. DNA sequences (+ indicates sense and -, antisense sequence) were read from the bottom to the top of each ladder in the directions indicated by the arrows (pointing to the right, 5' to 3' and to the left, 3' to 5'). The 87-bp exon is No. 5. Intron sizes were
15 determined by Southern blotting and restriction mapping and are accurate to within 10 percent, with one exception: intron 1 is accurate to within 20 percent.

20 (c) The structure of the human pituitary cDNA. Exon/intron splice junctions are indicated by triangles. Regions encoding transmembrane domains I-VII are enclosed by boxes containing wavy lines. The 87-bp sequence is striped. The sites for translation initiation (ATG) and termination (TAG) are indicated, as is the polyadenylation signal sequence (AATAAAA).

25 Figure 22 shows a comparison of IC_{50} values (nM) for L-hPitD2Zem, L-RGB2Zem1 and rat striatum.

30 Figure 23 shows the exon/intron junctions in the human dopamine D_2 receptor gene.

Discussion

35 This invention takes advantage in a unique way of nucleotide sequence similarities among members of a gene

family coding for receptors that are coupled to G proteins. By using a unique hamster β_2 -adrenergic receptor (β_2 AR) gene as a hybridization probe, a cDNA encoding the rat D_2 dopamine receptor was identified and isolated. The receptor has been characterized on the basis of three criteria: 1) the deduced amino acid sequence which reveals that it is a member of the family of G protein-coupled receptors, 2) the tissue distribution of the corresponding mRNA which parallels that known for the D_2 dopamine receptor, and 3) the pharmacological profile of Ltk- cells transformed with the cDNA.

A rat genomic library was screened under low-stringency hybridization conditions with a nick-translated 1.3 kb HindIII fragment containing most of the coding region of the hamster β_2 AR gene. The hamster β_2 AR receptor gene was cloned from a partial hamster lung λ gt10 genomic DNA library (constructed from size fractionated (5-7 kb) EcoRI digested DNA) with two oligonucleotide probes (30-mer, TCTGCTTTCAATCCCCTCATCTACTGTCGG; 40-mer, CTATCTTCTGGAGCTGCCTTTTGGCCACCTGGAAGACCCT) designed from the sequence of Dixon et al. (9). The 1.3 kb HindIII fragment of the hamster β_2 AR gene which contains most of the coding sequence of that gene was labeled by nick translation and used to probe a rat genomic DNA library in the commercially available phage EMBL3. The library was transferred to Colony Plaque Screen filters (NEN) and screened with the 32 P labeled probe using the following hybridization conditions: 25% formamide, 5 x SSC, 5 x Denhardtts, 0.1% sodium pyrophosphate, 1% SDS and salmon sperm DNA (100 μ g/ml) at 37°C. Filters were washed in 2 x SSC and 0.1% SDS at 55°C.

Several clones were found to hybridize to the hamster probe using these conditions. One clone, called RGB-2, was found to have a 0.8 kb EcoRI-PstI fragment that hybridized to the hamster β_2 AR probe in Southern blot

analysis. This fragment was sequenced and shown to have a stretch of nucleotides with a high degree of identity (32 out of 40 bases) to the nucleotide sequence of transmembrane domain VII of the hamster β_2 AR. One of the possible reading frames demonstrated a significant similarity to the amino acid sequence of transmembrane domains VI and VII of the hamster β_2 AR. Within this genomic fragment there is also a 3'intron splice site (10) and 400 bp of putative intronic sequence.

The 0.8 kb EcoRI-PstI fragment (nick translated) was used to probe a rat brain cDNA library in λ gt10 with the same hybridization conditions as above except that 50% formamide was used. Washing of the filters was performed in 0.2 x SSC and 0.1% SDS at 65°C. Under these high stringency hybridization conditions, two positive clones of about 2.5 kb in size were identified from a library of 500,000 clones. DNA sequence was obtained from both strands by the Sanger dideoxy chain determination method using Sequenase (U.S. Biochemical) (26). Sequence and restriction analysis indicated that the two clones were replicas of a single clone and contained the sequence of the genomic fragment that had been used as a probe. When the RGB-2 cDNA was used as a hybridization probe in Northern blot analysis of mRNA isolated from rat brain, a band of approximately 2.5 kb was detected. This finding indicated that the RGB-2 clone was nearly full length.

Figure 1 shows the nucleotide sequence of 2455 bases for the RGB-2 cDNA. The longest open reading frame in this cDNA codes for a 415 amino acid protein (relative molecular weight (M_r =47,064)) also shown in the figure. This molecular weight is similar to that reported for the deglycosylated form of the D_2 dopamine receptor as determined by SDS polyacrylamide gel electrophoresis (11). An in-frame dipeptide which is 36 bases upstream from the putative initiation site is found in the 5' untranslated region of the RGB-2 cDNA. A small open

reading frame has been observed in the 5' untranslated sequence of the β_2 AR mRNA (9).

Several structural features of the protein deduced from the RGB-2 cDNA demonstrate that it belongs to the family of G protein-coupled receptors. The hydrophobicity plot of the protein sequence (Figure 5) shows the existence of seven stretches of hydrophobic amino acids which could represent seven transmembrane domains (8). Moreover, the primary amino acid sequence of RGB-2 shows a high degree of similarity with other G protein-coupled receptors (Figure 2). The regions of greatest amino acid identity are clustered within the putative transmembrane domains. Within these domains the RGB-2 protein has a sequence identity of 50% with the human α_2 -adrenergic receptor (12), 42% with the human G-21 receptor (13), 38% with the hamster β_2 AR-adrenergic receptor (9), 27% with the porcine M_1 receptor (14), and 25% with the bovine substance K-receptor (15).

Thirdly, RGB-2 has several structural characteristics common to the members of the family of G protein-coupled receptors. There are three consensus sequences for N-linked glycosylation in the N-terminus with no signal sequence. The Asp 80 found in transmembrane domain II is conserved in all known G protein-coupled receptors. In transmembrane domain III there is Asp 114 for which a corresponding Asp residue is found only in receptors that bind cationic amines (16). Phosphorylation has been proposed as a means of regulating receptor function (17). A potential site for phosphorylation by protein kinase A exists at Ser 228 in the third cytoplasmic loop. In the C-terminal portion of the rhodopsins and β -adrenergic receptors are found many Ser and Thr residues which are potential substrates for receptor kinases (18). In contrast, the short C-terminus of RGB-2 has no Ser and Thr residues. However, as is the case for the α_2 -adrenergic receptor, RGB-2 has many Ser and Thr residues (22 residues) in the third cytoplasmic

loop which could serve as phosphorylation substrates. RGB-2 contains a large cytoplasmic loop (135 amino acids) between transmembrane domains V and VI with a short C-terminus (14 amino acids). This structural organization is similar to other receptors which are coupled by G_i (inhibitory G protein) such as the α_2 -adrenergic receptor and the M₂ muscarinic receptor. Unlike the members of the adrenergic and muscarinic receptor families, the RGB-2 gene has at least one intron in its coding sequence which is located in transmembrane domain VI.

As a first step towards determining the identity of RGB-2, the tissue distribution of the RGB-2 mRNA was examined by Northern blot analysis (Figure 3).

Brain tissue was dissected from male Sprague Dawley rats and RNA isolated according to Chirgwin et al. (27) and Ullrich et al. (28). For Northern blotting, RNA was denatured using glyoxal and DMSO and run on 1.2% agarose gels. After electrophoresis, RNA was blotted onto a nylon membrane (N-Bond, Amersham) and baked for 2 hours at 80°C. The membranes were prehybridized in 50% formamide, 0.2% PVP (M.W. 40,000), 0.2% ficoll (M.W. 400,000), 0.05 M Tris pH 7.5, 1M NaCl, 0.1% PPi, 1% SDS and denatured salmon sperm DNA (100µg/ml) for 16 hrs at 42°C. A random primed ³²P-labeled fragment (1-2 10⁸ dpm/µg) from the 1.6 kb BamHI-BglIII fragment of RGB-2 which contains the coding region of this clone was used at 10⁷ dpm/ml in the hybridization solution from above to probe the filters overnight at 42°C. The blots were washed twice in 2x SSC and 0.1% SDS at 65°C for 10 min., twice in 0.5x SSC and 0.1% SDS at RT for 15 min. and once in 0.1x SSC and 0.1% SDS at 65°C for 15 min. Blots were exposed overnight at -70°C to X-ray film with an intensifying screen.

The RGB-2 mRNA is expressed at different levels in various regions of the rat brain with the basal ganglia showing the highest concentration. Furthermore, the RGB-2 mRNA was found in high amounts in the neurointermediate

lobe of the pituitary gland of the rat and to a lesser degree in the anterior lobe of this gland. The expression pattern of the RGB-2 mRNA is strikingly similar to the distribution of the D₂ dopamine receptor as determined by receptor autoradiography and binding studies of tissue preparations (19).

In order to study the pharmacological characteristics of the receptor encoded by RGB-2, the cDNA was expressed in eucaryotic cells. The full RGB-2 cDNA was cloned into the eucaryotic expression vector pZem3 (20) which initiates transcription from the mouse metallothionein promoter (21). This plasmid was cotransfected with the selectable neomycin phosphotransferase gene (pRSVneo) into the Ltk- mouse fibroblast cell line by the standard CaPO₄ precipitation technique (21). Cells were selected in 350 µg/ml of G418. Transfectants were isolated and checked for expression of RGB-2 mRNA by Northern blot analysis. A cell line expressing RGB-2 designated L-RGB2Zem-1 was isolated. The RGB-2 mRNA was not detectable in the parent Ltk- cell line.

Since the RGB-2 mRNA displayed the tissue distribution expected of the D₂ dopamine receptor, a pharmacological study was performed of the L-RGB2Zem-1 cell line, native Ltk- cells and rat striatum using the D₂ ligand ³H-spiperone.

Membranes were prepared by homogenizing cells with a Dounce homogenizer at 4°C in 0.25 M sucrose, 25 mM Tris pH 7.4, 6 mM MgCl₂, 1 mM EDTA. The homogenizing solution was centrifuged at 800 x g for 10 min. and the pellet was subjected to a second homogenization and centrifugation as before. The supernatants were pooled and centrifuged at 200,000 x g for 1 hour. The pellet of this centrifugation was resuspended in 25 mM Tris pH 7.4, 6 mM MgCl₂, 1mM EDTA at approximately 250 µg protein/ml and stored in small aliquots at -70°C. Radioligand binding assays were carried out in duplicate in a volume of 2 ml (saturation analyses) or 1 ml (inhibition curves)

containing (final concentration): 50 mM Tris, pH 7.4, 0.9% NaCl, 0.025% ascorbic acid, 0.001% bovine serum albumin, ^3H -spiperone (Amersham, 95 Ci/mmol) and appropriate drugs. In some experiments 100 μM guanosine 5'-triphosphate was included. (+)-Butaclamol (2 μM) was used to define nonspecific binding. Incubations were initiated by the addition of 15-40 μg of protein, carried out at 37°C for 50 minutes, and stopped by the addition of 10 ml of ice-cold wash buffer (10 mM Tris, pH = 7.4, and 0.9% NaCl) to each assay. The samples were filtered through glass-fiber filters (Schleicher and Schuell No. 30) and washed with an additional 10 ml of wash buffer. The radioactivity retained on the filter was counted using a Beckman LS 1701 scintillation counter. Data were analyzed as previously described (29) except that curves were drawn using the data analysis program Enzfitter. The resulting IC_{50} values were converted to K_i values by the method of Cheng and Prusoff (30).

Membranes prepared from control Ltk- cells showed no (+)-butaclamol- or sulpiride-displaceable binding of ^3H -spiperone. Binding of ^3H -spiperone to membranes prepared from L-RGB2Zem-1 cells was saturable with a K_d value of 48 pM (Figure 4a). This value agrees with that observed for binding of ^3H -spiperone to rat striatal membranes in parallel experiments (52 pM). In the experiment shown in Figure 4a, K_d and B_{max} values for membranes prepared from L-RGB2Zem-1 were 40 pM and 876 fmol/mg of protein, whereas the corresponding values in striatal membranes were 37 pM and 547 fmol/mg of protein. The density of binding sites as determined in four experiments was 945 fmol/mg of protein in L-RGB2Zem-1 membranes and 454 fmol/mg of protein in rat striatal membranes.

The binding of ^3H -spiperone to membranes from L-RGB2Zem-1 cells was inhibited by a number of drugs and the resulting K_i values closely matched those obtained using striatal membranes (Figures 4b, 4c). The D_2 antagonists (+)-butaclamol and haloperidol were the most

potent inhibitors, followed by sulpiride. The D₁ dopamine antagonist SCH 23390 and the serotonin 5HT₂ antagonist ketanserin were much less potent at blocking ³H-spiperone binding. The binding appeared to be stereoselective as (+)-butaclamol was much more potent than (-)-butaclamol at inhibiting binding. In these experiments the absolute affinities of dopaminergic antagonists and the rank order of potency of the drugs (spiperone > (+)-butaclamol > haloperidol > sulpiride >> (-)-butaclamol) agree closely with previously published values for the D₂ dopamine receptor (2).

All binding data for L-RGB2Zem-1 membranes were fit best by assuming the presence of only one class of binding sites. On the other hand, inhibition by several drugs of ³H-spiperone binding to rat striatal membranes was fit best by assuming the presence of two classes of binding sites. Thus, SCH 23390 and ketanserin inhibited 10-20% of ³H-spiperone binding to rat striatal membranes with high affinity (Figure 4c). In rat striatal membranes, inhibition of radioligand binding by sulpiride was fit best by one class of binding sites, but 10-15% of the (+)-butaclamol-displaceable binding was not inhibited by sulpiride at the concentrations used. It seems likely that the binding sites with high affinity for ketanserin and SCH 23390 and which are not displaced by sulpiride represent binding of ³H-spiperone to 5HT₂ serotonin receptors in rat striatal membranes. Binding of SCH 23390 to 5HT₂ receptors has been described previously (22). In rat striatal membranes, the apparent affinity of drugs for D₂ dopamine receptors, the class of binding sites comprising 80-90% of ³H-spiperone binding, was indistinguishable from the apparent affinity of drugs to membranes prepared from L-RGB2Zem-1 cells (Figure 4c).

The physiological effects of stimulation of D₂ dopamine receptors appear to be mediated by G_i (6). Inhibition of agonist binding to D₂ dopamine receptors by GTP is thought to be due to GTP-induced dissociation of a

receptor-G_i complex which has a high affinity for agonist (23,24). Although ³H-spiperone binding was inhibited by the agonist dopamine with a K_i of 17 μ m in the L-RGB2Zem-1 membranes, this dopamine binding was not responsive to the addition of GTP. However, this finding is consistent with the reported lack of G_i in L cells (25). The pharmacological data presented here proves that the binding profile of the D₂ dopamine receptor is found in Ltk- cells expressing the RGB-2 cDNA.

The foregoing data show that, when transfected into eucaryotic cells, the RGB-2 cDNA directs the expression of a D₂ dopamine binding protein. Since the mRNA corresponding to this cDNA is localized in tissues where the D₂ dopamine receptor is known to be present and since this mRNA codes for a protein which has all the expected characteristics of a G protein-coupled receptor, inter alia, RGB-2 is a clone for the rat D₂ dopamine receptor.

The nucleic acid sequence shown in Figure 1 can be inserted into a wide variety of conventional and preferably commercially available plasmids, e.g., using EcoRI sites or other appropriate sites. See, e.g., Figure 6 for a restriction map of the sequence of Figure 1.

Dopamine receptor genes of this invention, particularly mammalian D₂ dopamine receptor genes, based on this disclosure, can now be routinely made, isolated and/or cloned, using many conventional techniques. For example, the procedure disclosed herein can be substantially reproduced for libraries containing dopamine receptor DNA sequences. Alternatively, oligonucleotide probes can be routinely designed, e.g., from the sequences of Figure 1 and/or the omitted introns, which are selective for dopamine receptor genes, especially for mammalian dopamine D₂ receptor genes. These can be used to screen nucleic acid libraries containing dopamine receptor nucleic acid sequences. Sequences in these libraries hybridizing to the probes,

especially to all of a plurality of such probes (e.g. 2 or 3), will be DNA sequences of this invention with high probability. Of course, it is also possible to synthesize the sequence of Figure 1 or any fragment thereof using conventional methods.

This invention also enables the production of a wide variety of useful products and the employment of a wide variety of useful methods, as well as providing basic tools for the study of the regulation and function of dopamine receptors.

These products and methods can be produced and carried out, respectively, using the well known recombinant DNA, immunochemical and other methodologies of the biotech industry. See, e.g., USP 4,237,224; USP 4,264,731; USP 4,273,875; USP 4,293,652; EP 093,619; Davis et al "A Manual for Genetic Engineering, Advanced Bacterial Genetics," Cold Spring Harbor Laboratory, Cold Spring Harbor, New York (1980); Maniatis et al., Molecular Cloning, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.; Davis et al., Basic Methods in Molecular Biology, Elsevier, N.Y. (1986); Methods in Enzymology, Berger & Kimmel (Eds.), (1987).

This invention provides, for the first time, purified and isolated polypeptide products having all or part of the primary amino acid sequence of the dopamine D₂ receptor as well as its biological properties, including: all sites for covalent modification, such as phosphorylation and glycosylation; all primary sequence that determines the secondary, tertiary and quaternary structure of the functional protein; all parts of the molecule that provide antigenicity and antibody binding sites; all parts of the protein that provide for noncovalent interactions with other biological molecules, such as lipids, carbohydrates, and proteins, such as guanine nucleotide-binding regulatory proteins; all parts of the protein that make up the binding sites for ligands, including agonists, partial agonists and

antagonists; all parts of the protein that are required for functional conformational changes involved in normal biological activities, such as desensitization; and all proteins that might arise as a result of alternative splicing of the gene for this receptor.

These polypeptides can be expressed from the nucleic acids of this invention by procaryotic or eucaryotic hosts, e.g., bacterial, yeast or mammalian cells in culture, using fully conventional transformation or transfection (e.g., via calcium phosphate for mammalian cells) techniques. The products of such expression in vertebrate (e.g., mammalian and avian) cells are especially advantageous in that they are produced free from association with other human proteins or contaminants with which they may be associated in natural form. Preferred hosts for expression are mammalian and include for example mouse Ltk cells, hamster CHO cells, mouse GH₄ cells, mouse C₆ cells, mouse/rat NG108-15 cells and mouse AtT20 cells. For example, when the gene of Figure 1 is transfected into the commercially available growth hormone GH₄ cells, modulation of the cAMP second messenger system has been observed. Preferred vectors include pZem or pRSV or viral vectors such as vaccinia virus and retroviruses.

The polypeptides of this invention include all possible variants, e.g., both the glycosylated and non-glycosylated forms. The particular carbohydrates involved will depend on the mammalian or other eucaryotic cells used for the expression. The polypeptides of the invention can also include an initial methionine amino acid residue.

Also included in this invention are polypeptides, synthetic or otherwise, duplicating the amino acid sequence of Figure 1 and/or of the dopamine receptors per se of this invention, or only partially duplicating the same. These wholly or partially duplicative polypeptides will preferably also retain the biological and/or

immunological activity of the dopamine receptor per se. Also included within the scope of this invention are the monoclonal and polyclonal antibodies (generatable by conventional techniques and preferably labelled) which
5 are immunoreactive with such polypeptides.

Preferred partial polypeptides (fragments) are those including at least a portion of the sequences located in the hydrophobic transmembrane domains V, VI and VII, shown in Figure 2. These are the likely locations of the
10 ligand binding site(s), particularly domain VII. The third cytoplasmic loop is also an important fragment area; e.g., G-protein binding requires this location as well as domains V and VI. Where it is desired to have an antibody highly specific to a particular dopamine
15 receptor, a fragment generating such an antibody will be selected from the highly unique region between transmembrane regions V and VI, i.e., the third cytoplasmic loop, or the C-terminal domain, both of which have low homology with other receptors, and/or the
20 antibodies will be selected to be specific to an epitope in these regions. Another receptor/gene specific region is that of the intron sequences, e.g., those for RGB-2, mentioned above. Particularly preferred peptides which have been synthesized are (referring to the amino acid
25 numbers of Figure 1): (A) 2-13; (B) 182-192; (C) 264-277; (D) 287-298; and (E) 404-414. These are selected based on the following principles: the known antigenicity of peptides containing a large number of Pro residues (B/C/D); coverage of the N and C termini (A) and
30 (E); the ability to direct antibodies towards an extracellular domain (receptor reaction region) which will be effective to block the receptor reaction (A/C/D). Antibodies to these fragments are raised conventionally, e.g., monoclonals by fully conventional hybridoma
35 techniques.

This invention also relates to DNA sequences encoding the full dopamine receptor or fragments thereof,

as well as expression vectors (e.g., viral and circular
plasmid vectors) containing such whole or partial
sequences. Similarly, hosts (e.g., bacterial, yeast and
mammalian) or cells transformed or transfected with such
5 vectors are also included. The corresponding methods of
expressing the polypeptides corresponding to the
sequences in such vectors are also included, e.g.,
comprising culturing the transformed or transfected cells
under appropriate conditions for large scale expression
10 of the exogenous sequences and for the isolation of the
polypeptides as usual, e.g., from the growth medium,
cellular lysates or cellular membrane fractions.

DNA sequences of this invention also include those
coding for polypeptide variants, mutants or analogues
15 which differ from the natural sequence described herein.
Such differences can be derived from deletion of one or
more amino acid residues from the natural sequence, by
substitution of a given such residue by another residue
and/or by additions wherein one or more amino acid
20 residues are added to the natural sequence. Preferably,
the resultant modified polypeptide will retain at least
one of the biological or immunological activities of the
dopamine receptor.

Mutations likely to affect dopamine affinity acti-
25 vity will be those in transmembrane domains V, VI or VII
or in the third cytoplasmic loop. In addition, the DNA
sequences also include sequences complementary to any of
the other DNA strands mentioned herein and, most notably,
those shown in the Figures; DNA sequences which hybrid-
30 ize to the DNA sequences described herein, typically
under the hybridization conditions mentioned herein or
under more stringent conditions, or which hybridize to
fragments of such DNA sequences; and DNA sequences which
differ from those shown herein by the degeneracy of the
35 genetic code. Thus, this invention includes all DNA se-
quences which encode a dopamine receptor and hybridize to
one or more of the sequences shown herein. These include

allelic variants as well as dopamine receptors from mammalian species other than the species mentioned in the experimental descriptions herein.

5 Modifications of the cDNA or genomic dopamine receptor DNA may be readily accomplished by any of the well known techniques, including site-directed mutagenesis techniques. Such modified DNA sequences can include deletions, additions and/or substitutions made in selected regions, e.g., not in transmembrane domains V,
10 VI or VII or in the third cytoplasmic loop, where retention of the underlying biological activity of the dopamine receptor is desired.

Modified proteins which do not retain the mentioned biological activity and/or the corresponding DNA
15 sequences will also be useful, e.g., in various assays of this invention. In a particularly preferred such modification, the transmembrane domain V, VI, or VII or the third cytoplasmic loop will be deleted or rendered inactive, e.g., by sequence modification. Deletion of
20 the glycosylation sites shown in Figure 1 is also a useful variant for expression of the polypeptide, e.g., in yeast cells.

As mentioned above, it is well established that significant portions of the DNA sequence encoding a
25 dopamine receptor are conserved in various mammalian species. Consequently, using only routine experimentation, a skilled worker can readily screen a DNA genomic library or, preferably, a cDNA library, e.g., from the brain of a given mammal, for the presence of other
30 dopamine receptor genes, especially D₂ dopamine receptor genes, using probes manufactured in accordance with the details of the sequences shown herein, including the 5' flanking, the intronic and the structural gene sequences shown in Figure 1 and the human sequence of Figure 7.
35 Probes will preferably be selected from the seven highly conserved transmembrane domains shown in Figure 2, preferably domains VI and VII. Such a routine screening

will identify clones which hybridize with the probes. From these, dopamine receptors can routinely be selected, e.g., using the techniques described herein. With respect to human D₂ dopamine receptors, particularly

5 useful sources include, for cDNA, striatum, pituitary, neuroblastoma, kidney, placenta cells, etc., and, for genomic DNA, liver, placenta cells, etc. For primates, e.g., rhesus monkeys, particularly useful genomic DNA or cDNA libraries include brain, kidney and placenta cells.

10 With respect to human D₂ dopamine receptor genes, the partial sequence shown in Figure 7 has been identified by conventionally screening, under the stringent hybridization conditions described above for the probing by the 0.8 kb EcoRI-PstI fragment of rat brain cDNA in

15 λ gt10, a pituitary cDNA library using a probe which is the full length rat cDNA, RGB-2. The cDNA libraries mentioned herein were prepared by fully conventional methods, e.g., as described in the references cited above, e.g., Davis et al. This sequence or fragments thereof can also be useful as a probe, for example, to

20 screen conventional libraries as mentioned above for human dopamine receptor genes in accordance with the foregoing and other fully conventional procedures. As can be seen by comparing the sequence of Figure 7 with

25 the sequences shown in Figures 1 and 2 above, the partial human sequence of Figure 7 has high homology with RGB-2 beginning at amino acid no. 259 of Figure 1.

Similarly, this invention more generally includes mammalian D₂ dopamine receptor genes in the broadest

30 sense, e.g., both regulatory and structural such genes, alone or in combination, e.g., in reading frame. For example, using the routine methods discussed herein, such genes have been and can be cloned from mammalian DNA libraries. As well, this invention includes biologically

35 active fragments of such genes, e.g., fragments encoding polypeptides having the biological activity of a mammalian D₂ dopamine receptor, or fragments useful in

controlling expression of such encoding fragments, or fragments useful as probes for any such gene or fragment, e.g., by hybridizing therewith.

5 The various polypeptides and sequences of this invention may be conventionally labeled with detectable marker substances, typically by covalent association, and further typically by radiolabeling, or in the case of DNA, with non-isotopic labels such as biotin. The polypeptide products (e.g., labelled antibodies) can be
10 used conventionally to detect and quantitate the presence of dopamine receptors in various samples; the DNA-labeled products can be conventionally employed in the usual hybridization methods (e.g., Northern blots, Southern blots, spot assays, etc.) to detect and quantitate the
15 presence of associated nucleic acids (DNA, RNA) in samples, e.g., to locate the dopamine gene positions in various mammalian chromosomal maps, to determine whether mRNA or receptor concentrations are abnormally high or low in comparison to standard levels, etc. They will
20 also be useful, again using fully conventional procedures, to identify dopamine receptor gene disorders (defective or aberrant genes) in in vitro diagnostic procedures on DNA samples from given patients, e.g., to detect chromosomal defects, e.g., using RFLP analyses
25 (see, e.g., Genes III, Levin, John Wiley and Sons (1987)). For example, since dopamine receptors have been implicated in schizophrenia, products and methods of this invention can be used to characterize nucleic acids, e.g., in size fractionated form, from schizophrenia
30 patients and, inter alia, classify patients according to schizophrenia subgroups, make diagnoses based on comparisons to standard DNA fractionation arrays, etc. Of course, they can also be used, e.g., in the mapping of the human or other mammalian genome, as gene markers to
35 identify accompanying genes, e.g., in RFLP analyses, and, where applicable, other disorders. The gene-unique regions discussed above, e.g., the intron regions, the

third cytoplasmic loop, etc., will be especially useful in this regard.

Typical assays in which the polypeptides of this invention can be utilized include any of the well known
5 immunoassay techniques such as RIA, ELISA, etc., both of in vitro and in vivo nature. Various fragments of the polypeptide sequence of the dopamine receptor can also be utilized conventionally for producing corresponding polyclonal antibodies or preferably monoclonal antibodies
10 (e.g., by conventional preparation and expression of corresponding hybridomas) for epitopes within the given fragment. The antibodies will often be conventionally labelled, e.g., radio- or enzymatically labelled. In a preferred aspect, the resultant polyclonal or monoclonal
15 antibodies will also be immunospecific with respect to not only the mentioned fragment, but also the full protein. Such antibodies will be conventionally employable, for example, in the detection and affinity purification or chromatography of dopamine receptor and
20 related products.

Of course, the polypeptides of this invention include those expressed in accordance with conventional procedures from cells as mentioned above, as well as those which are synthetically prepared also using
25 conventional procedures. This invention enables for the first time non-natural preparation of mammalian dopamine receptors substantially free of constituents of their natural environment. It is in this sense the term "substantially pure" is used herein, i.e., substantially
30 free of these natural constituents. The invention also includes DNA sequences which can be isolated from the various sources mentioned above or synthetically prepared using fully conventional methods.

Also included within the scope of this invention are
35 pharmaceutical compositions comprising effective amounts of one or more of the polypeptide products of this invention or one or more of the nucleic acid sequences of

this invention, in admixture with suitable conventional diluents, adjuvants and/or carriers well known in the pharmaceutical industry. These can be utilized for in vitro uses, e.g., for detection of the presence of a dopamine receptor in a sample or of the presence of a gene or an abnormal gene in a sample or for increasing the concentration of receptor or its gene in a sample, or for in vivo uses such as gene therapy (e.g., to render a defective gene or gene product inactive, e.g., block it by an appropriate monoclonal antibody) and/or to provide new functional genes (e.g., using retroviral vectors)), or to provide an increased concentration of dopamine receptor in a given location, or to modulate receptor expression and/or activity, e.g., by administration of antisense oligosequences, all in mammals including humans. Thus, these compositions will be useful to treat disease conditions, inter alia, those associated with abnormalities in the structure, expression or concentration of the dopamine receptor or its gene, such as those mentioned in the foregoing. Specific effective dosages for a given condition in a given patient will vary, as is well known, with the usual conditions, including the overall condition of the patient, body weight, the identity and severity of the particular dopamine-deficiency disease state, etc.

The polypeptides of this invention can also be used in, e.g., competitive binding assays, to test for the affinity thereto of candidate chemical substances such as drugs, e.g., affinity (e.g., agonistic or antagonistic) to D₂ dopamine receptors. Such procedures can be carried out, e.g., as pharmaceutical screening tests, using fully conventional procedures, analogous to those described herein and/or to known protocols based on natural sources of dopamine receptors, e.g., analogous to known tests for inhibition of the binding of tritiated dopamine agonists and antagonists to striatal receptors per the methods of Schwarcz et al., J. Neurochemistry, 34 (1980), 772-778

and Creese et al., European J. Pharmacol., 46 (1977),
377-381, and to those for other receptors. It is also
possible to screen substances for ability to modify or
initiate a response which is triggered by ligand bonding
to a dopamine receptor, e.g., cellular responses such as
modulation of second messenger systems. Such analyses
can utilize cells of this invention transformed with
nucleic acid sequences of this invention.

Antibodies of this invention to the dopamine
receptors or other regions of dopamine receptor genes,
especially the D₂ receptor, can also be used in diagnostic
imaging techniques, e.g., by radiodiagnostic, MRI or
positron imaging. Radio-, paramagnetic- or positron-
labels can be conventionally attached to the antibodies
(preferably monoclonal in nature), e.g., via covalent
bonding to chelating groups for a positron emitting,
radionucleotide or paramagnetic metal. MRI,
radiosensitive or positron imaging can then be effected
with these agents using conventional methods. See, e.g.,
Maziere et al., Life Sci., 35, 1349 (1984).

Suitable pharmaceutical carriers include water,
saline, human serum albumin, etc. The compositions can
also include other active ingredients suitable for
amelioration of the particular disease state involved,
e.g., conventional dopamine agonists, dopamine
antagonists, etc. The components of this invention can
be provided in conventional kit form containing, e.g., an
antibody or a DNA probe (e.g., able to detect gene
homologies or anomalies) along with detection method-
specific reagents such as enzymes, substrates, materials
for analyzing DNA restriction fragments, etc.

The DNA sequences of this invention are also useful
to prepare the corresponding transgenic animals, in
particular nonhuman mammals, e.g., rats, monkeys, etc.,
using known methods, e.g., analogous to those described
in USP 4,736,866. Such animals, e.g., are particularly
useful for commercial research purposes. The DNA

sequences or the corresponding mRNA can also be used conventionally to inject oocytes, e.g., from frogs, which can then be conventionally used in binding or second messenger analyses. Moreover, the availability of the primary amino acid sequence itself enables experimental and computational modeling and understanding of the secondary and tertiary structures of the dopamine receptor. This three-dimensional information provides a basis for modeling and understanding the details of the receptor function, e.g., interaction with the cell membrane, its ligand (binding pocket), associated proteins, messenger systems, etc. Such analyses enable rational drug design whereby, e.g., new dopamine receptor affecting chemical agents can be designed in accordance with the details of this interaction.

Numbered References for Background, Summary,
Figures 1-7, and Discussion (Excluding Examples)

1. Creese, I., Sibley, D.R., Hamblin, M.W. & Leff,
S.E. Ann. Rev. Neurosci. 6, 43-71 (1983).
- 5 2. Seeman, P. Pharmacol. Rev. 32, 229-313 (1980).
3. Lee, T. et al. Nature 273, 59-61 (1984).
4. Seeman, P. et al. Science 225, 728-731 (1984).
5. Barnes, D.M. Science 241, 415-417 (1988).
6. Cote, T.E., Frey, E.A., Grewe, C.W., & Keabadian, J.W.
10 J. Neural. Trans. Suppl. 18, 139-147 (1983).
7. Senogles, S.E. et al. J. Biol. Chem. 262,
4860-4867 (1987).
8. Dohlman, H.G., Caron, M.G. & Lefkowitz, R.J.
Biochemistry 26, 2657-2664 (1987).
- 15 9. Dixon, R.A.F., et al. Nature 321, 75-79 (1986).
10. Mount, S.M. Nucleic Acids Res. 10, 459-472 (1982).
11. Grigoriadis, D.E., Niznik, H.B., Jarvie, K.R. &
Seeman, P. FEBS Let. 227, 220-224 (1988).
12. Kobilka, B.K., et al. Science 238, 650-656 (1987).
- 20 13. Kobilka, B.K., et al. Nature 329, 75-79 (1987).
14. Kubo, T., et al. Nature 323, 411-416 (1986).
15. Kubo, Y., et al. Nature 329, 836-838 (1986).
16. Strader, C.D., et al. J. Biol. Chem. 263, 10267-
10271 (1988).
- 25 17. Sibley, D.R., Benovic, J.L., Caron, M.G. & Lefkowitz,
R.J. Cell 48, 913-922 (1987).
18. Bouvier, M., et al. Nature 333, 370-373 (1988).
19. Boyson, S.J., McGonigle, P. & Molinoff, P.B. J.
Neurosci. 6, 3177-3188 (1986).
- 30 20. Uhler, M. & McKnight, G.S. J. Biol. Chem. 262,
15202-15207 (1987).

21. Gorman, C., Padmanabhan, R. & Howard, B.H. Science 221, 551-553 (1983).
22. Hyttel, J. Eur. J. Pharmacol. 91, 153-154 (1983).
23. Hamblin, M.W., Leff, S.E. & Creese, I. Biochem. Pharmacol. 33, 872-877 (1984).
24. Dolphin, A.C. Trends in Neurosci. 10, 53-57 (1987).
25. Jones, S.V.P., et al. Proc. Nat. Acad. Sci. U.S.A. 85, 4056-4060 (1988).
26. Sanger, F., Nicklen, S. & Coulson, A.R. Proc. Nat. Acad. Sci. U.S.A. 74, 5463-5467 (1977).
27. Chirgwin, J.M., Przybyla, A.E., McDonald, R.J. & Rutter, W.J. Biochemistry 18, 5294-5299 (1979).
28. Ullrich, A., et al. Science 196, 1313-1319 (1977).
29. Neve, K.A. & Molinoff, P.B. Mol. Pharmacol. 30, 104-111 (1986).
30. Cheng, Y.C. & Prusoff, W.H. Biochem. Pharmacol. 22, 3099-3108 (1973).

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The mentioned embodiments are, therefore, to be construed as
5 merely illustrative and not limitative of the remainder of the disclosure in any way whatsoever.

From the foregoing description, one skilled in the art can easily ascertain the essential characteristics of this invention and, without departing from the spirit and
10 scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions.

The entire texts of all applications, patents and publications cited above are hereby incorporated by
15 reference.

EXAMPLES

Example 1

Summary

We recently cloned a complementary DNA for the rat
20 dopamine D-2 receptor, making it possible to create cell lines expressing this receptor. A cell line (LZR1) was created by transfecting the D-2 cDNA (RGB-2) into mouse fibroblast Ltk⁻ cells. LZR1 cells, previously described as L-RGB2Zem-1 cells (1), expressed a high density of
25 D-2 receptors, whereas the wild-type cells did not. Although transcription of the RGB-2 cDNA is regulated by the zinc-inducible mouse metallothionein promoter, the density of D-2 receptors on membranes prepared from LZR1 cells was not increased after the cells were treated with
30 zinc. A second cell line derived from Ltk⁻ cells, designated LZR2, had a lower density of D-2 receptors in the uninduced state, and treatment with zinc induced a 50% increase in the receptor density. A number of agonists competitively and stereoselectively inhibited
35 the binding of [³H]spiroperidol to the expressed D-2 receptors. In LZR1 cells, dopamine was a more potent

inhibitor of radioligand binding in the absence than in the presence of GTP and NaCl. Dopamine reduced forskolin-stimulated adenylate cyclase activity by 27% in membranes prepared from LZR1 cells. Inhibition by dopamine was blocked by (+)-butaclamol or prior treatment of intact cells with pertussis toxin. These data indicate that the RGB-2 cDNA directs the expression of a dopamine D-2 receptor capable of interacting with guanine nucleotide-binding proteins and inhibiting adenylate cyclase activity. Furthermore, the RGB-2 cDNA provides a means of creating many cell lines that will be useful tools for the biochemical and pharmacological characterization of dopamine D-2 receptors.

Introduction

Dopamine (DA) receptors have been classified into two subtypes based on functional and pharmacological profiles (2). DA D-2 receptors are characterized functionally by their ability to inhibit adenylate cyclase activity (3). Activation of D-2 receptors also inhibits calcium channels (4,5), increases potassium conductance (6), and may inhibit accumulation of inositol phosphates (7,8). One factor that has impeded research on the regulation and functional characteristics of DA receptors has been the lack of cell lines that express the receptors. One cell line, derived from a prolactin-secreting tumor, has recently been described in which DA inhibits adenylate cyclase activity and prolactin secretion (9).

We recently cloned a rat brain complementary DNA (cDNA), designated RGB-2, that has significant homology with β_2 -adrenergic receptors and other receptors that interact with guanine nucleotide-binding proteins. Three lines of evidence indicate that the RGB-2 cDNA encodes the DA D-2 receptor: (1) The deduced amino acid-sequence of the protein suggests the existence of the seven membrane-spanning domains typical of receptors coupled to

guanine nucleotide-binding proteins (10); (2) the distribution of messenger RNA that hybridizes with the cDNA parallels the distribution of the D-2 receptor; and (3) when the RGB-2 cDNA is transfected into cells that lack high affinity binding of the D-2 selective ligand [³H]spiroperidol, the cells express binding sites for the radioligand with a pharmacological profile characteristic of D-2 receptors (1).

The cloning of a D-2 receptor cDNA makes it possible to express DA receptors in cell lines in which the effects of receptor activation can readily be determined. We previously described the binding of [³H]spiroperidol and other D-2 antagonists to a line of cells derived by transfection of mouse L cells with the RGB-2 cDNA under the control of a zinc-inducible mouse metallothionein promoter (1). We also reported that, under the assay conditions used previously, the binding of DA to LZR1 membranes was not responsive to GTP. We now demonstrate that the D-2 receptor encoded by the RGB-2 cDNA is functional with respect to the guanine nucleotide sensitivity of the binding of agonists and the ability of agonists to inhibit adenylate cyclase activity.

Methods

Materials: [α -³²P]Adenosine 5'-triphosphate (ATP, 10-50 Ci/mmol) and [³H]cyclic AMP (31.9 Ci/mmol) were purchased from New England Nuclear (Boston, MA), and [³H]spiroperidol (95 Ci/mmol) was purchased from Amersham (Arlington Heights, IL). Guanosine 5'-triphosphate (GTP), DA, cyclic AMP, 3-isobutyl-1-methyl-xanthine, ATP, and forskolin were purchased from Sigma Chemical Company (St. Louis, MO). Quinpirole, LY181990 (Lilly Laboratories), bromocriptine (Sandoz Research Institute), and (+) and (-)3-PPP (Astra) were generous donations.

Transfection: The full RGB-2 cDNA was cloned into the plasmid pZem3 (11). The cDNA and the vector were made compatible by partially filling in the Bgl II site

on the vector and a Sal I site on the cDNA adaptor. This plasmid was co-transfected with the plasmid pRSVneo into mouse Ltk⁻ cells by a CaPO₄ precipitation technique (12). Transfectants were selected in 350 µg/ml of G418, isolated, and screened for expression of RGB-2 mRNA by Northern blot analysis. The subclone LZR1, selected on the basis of high expression of RGB-2 mRNA, was partially characterized previously as L-RGB2Zem-1 (1). A second cell line, LZR2, was isolated in the same way.

Tissue culture: Cells were plated at a density of 20,000 cells/cm² in 150 mm diameter Falcon tissue culture plates (Beckton Dickinson, Lincoln Park, NJ), subcultured by replacing the growth medium with trypsin-EDTA (0.1% trypsin, 0.02% EDTA in phosphate-buffered saline) or fed on day 3, and harvested on day 5 or 6. Cells were grown in Dulbecco's modified Eagles' medium (Sigma), supplemented with 5% fetal bovine serum and 5% iron-supplemented calf bovine serum (Hyclone, Logan, UT), in an atmosphere of 10% CO₂/90% air at 37°. Cells were lysed by replacing the growth medium with ice-cold hypotonic buffer (1 mM Na⁺-HEPES, pH 7.4, 2 mM EDTA). After swelling for 10-15 min, the cells were scraped from the plate and centrifuged at 24,000 x g for 20 min. The resulting crude membrane fraction was resuspended with a Brinkmann Polytron homogenizer at setting 6 for 10 sec in Tris-isosaline (50 mM Tris-HCl, pH 7.4, and 0.9% NaCl) and stored at -70° for receptor binding experiments or resuspended in Tris-isosaline, centrifuged again at 24,000 x g for 20 min., and resuspended in Tris-isosaline for immediate use in adenylate cyclase experiments.

Receptor binding assay: The membrane preparation was thawed, centrifuged at 24,000 x g for 20 min., and resuspended in Tris-isosaline except where indicated. Aliquots of the membrane preparation were added to assay tubes containing (final concentrations) 50 mM Tris-HCl, pH 7.4, 0.9% NaCl, 0.025% ascorbic acid, 0.001% bovine serum albumin, [³H]spiroperidol, and appropriate drugs.

(+)-Butaclamol ($2 \mu\text{M}$) was used to define nonspecific binding, which was typically less than 10% of total binding at concentrations of radioligand near the K_D value. Assays were carried out in duplicate in a volume of 2 ml for saturation analyses or 1 ml for inhibition analyses. Incubations were initiated by the addition of 15-50 μg of protein, carried out at 37° for 50 min., and stopped by the addition of 10 ml of ice-cold wash buffer (10 mM Tris, pH 7.4, and 0.9% NaCl) to each assay. The samples were filtered through glass-fibre filters (Schleicher & Schuell No. 30) and washed with an additional 10 ml of wash buffer. The radioactivity retained on the filters was counted using a Beckman LS 1701 scintillation counter. Data were analyzed by nonlinear regression using the data analysis program Enzfitter (Elsevier-Biosoft). In competition experiments, K_i values were calculated from experimentally determined IC_{50} values by the method of Cheng and Prusoff (13). Averages for K_i and K_D values are the geometric means. In experiments designed to assess the effect of GTP and NaCl on the binding of DA, fresh tissue was used. Cells were harvested, centrifuged, and Mg^{2+} resuspended in Tris- Mg^{2+} (50 mM Tris, pH 7.4, and 4 mM MgCl_2). Tissue was incubated for 15 min. at 37° in this buffer before re-centrifugation. The resuspended protein was added to assays containing Tris- Mg^{2+} with no added NaCl or GTP, or Tris- Mg^{2+} with 120 mM NaCl and 100 μM GTP.

Adenylate cyclase assay: The conversion of $[\alpha^{32}\text{P}]\text{ATP}$ to $[\text{P}^{32}]\text{cAMP}$ was determined essentially as described by Salomon et al. (14). Membranes (50 - 100 μg of protein) resuspended in Tris-isosaline were added in a volume of 0.1 ml to an assay of 0.2 ml containing 50 mM Tris-HCl, pH 7.4, 5 mM cAMP, 1 mM 3-isobutyl-1-methylxanthine 1 mM MgCl_2 , 0.5 mM EGTA, 0.25 mM ATP, 30 μM GTP, approximately 2×10^6 cpm of $[\alpha^{32}\text{P}]\text{ATP}$, and various drugs. Assays were initiated by warming to 25° and terminated after 20 minutes by cooling to 0° , then adding

trichloroacetic acid (100 μ l of a 30% solution) to each assay. [3 H]Cyclic AMP (approximately 30,000 cpm) was added to each assay as an internal standard. The assay volume was brought up to 1 ml with water, and tubes were centrifuged for 10 min. at 2000 x g. Cyclic AMP in the supernatant was isolated by sequential chromatography on columns containing Dowex AG50W-X4 resin and neutral alumina. The 2-ml eluate from each column of alumina was dissolved in 10 ml of Bio-Safe II (RPI, Mount Prospect, IL) for liquid scintillation counting. Dose-response curves for inhibition of adenylate cyclase activity by agonists were analyzed by nonlinear regression using the program Enzfitter. The data were fit to the equation:

$$E = (100 - E_{\max}) / (1 + (A/EC_{50})^N) + E_{\max}$$

where E is the amount of enzyme activity, expressed as percentage of total stimulated activity, A is the concentration of agonists, EC_{50} is the concentration of agonist causing half-maximal inhibition of enzyme activity, E_{\max} is the enzyme activity observed in the presence of maximally inhibiting concentrations of agonist, expressed as the percentage of total stimulated activity, and N is a slope factor. Averages of EC_{50} values are the geometric means. Protein concentration was determined by the method of Peterson (15).

Results

Saturation analysis of the binding of

[3 H]spiroperidol: The density of D-2 receptors on membranes prepared from LZR1 cells was determined by saturation analysis of the binding of [3 H]spiroperidol (Fig. 8). Since the RGB-2 cDNA is under the control of the zinc-inducible mouse metallothionein promoter, the effect of prior treatment of cells with 100 μ M zinc sulfate on the binding of [3 H]spiroperidol was also determined. The density of binding sites was 736 ± 140 fmol/mg of protein in control LZR1 cells, and 759 ± 155 fmol/mg of protein in LZR1 cells treated with zinc (n =

2). A second cell line, designated LZR2, had a lower density of D-2 receptors (mean $B_{max} \pm SE$, 435 ± 71 fmol/mg of protein). In contrast to LZR1 cells, the density of binding sites on LZR2 cells was increased 50% by zinc treatment to 630 ± 52 fmol/mg of protein. The mean K_D value of 42 pM for control LZR1 and LZR2 cells ($pK_D \pm SE$, 10.37 ± 0.15 , $n = 4$) did not differ significantly from the mean K_D value of 47 pM for zinc-treated cells (10.33 ± 0.17). Scatchard transformation of saturation analyses from all experiments yielded straight lines. Wild-type Ltk⁻ cells had no detectable displaceable binding of [³H]spiroperidol (data not shown). Since LZR1 cells had the higher density of D-2 receptors, these cells were used in all subsequent experiments.

Inhibition of radioligand binding by agonists: The apparent affinity of D-2 receptors for several agonists and related compounds was determined in two experiments (Fig. 9A, Table 1). Of the six compounds tested, bromocriptine was the most potent with a mean K_i value of 2 nM, whereas LY181990, the inactive enantiomer of the D-2-selective agonist quinpirole, was the least potent. All assays were carried out in the presence of 0.1 mM GTP and 120 mM NaCl, using membranes prepared from LZR1 cells.

In other experiments, the effect of GTP and NaCl on inhibition of radioligand binding by DA was determined (Fig. 9B). Freshly prepared membranes were used for these experiments, and 4 mM $MgCl_2$ was added to the homogenization and assay buffers. In the absence of GTP and NaCl, the mean IC_{50} was 25 μM [mean $-\log(IC_{50}) \pm SE = 4.6 \pm 0.3$]. Addition of GTP and NaCl to the assay buffer increased the mean IC_{50} value to 167 μM (3.8 ± 0.3 , $n = 4$), without altering the binding of [³H]spiroperidol. Hill coefficients in the absence of GTP and NaCl ranged from 0.56 to 0.68 (mean = 0.64) whereas in the presence of GTP and NaCl values ranged from 0.77 to 1.1 (mean = 0.94). In the absence of GTP and NaCl, inhibition curves

for DA could be fit best by assuming the presence of two classes of binding sites. One class of high affinity sites, representing $48 \pm 14\%$ of the total number of receptors, had a mean K_i for DA of $0.3 \mu\text{M}$. The second class, representing $52 \pm 13\%$ of the total number of receptors had a mean K_i value of $24 \mu\text{M}$.

Inhibition of adenylate cyclase activity by agonists: In freshly prepared membranes from LZR1 cells, but not in membranes from wild-type Ltk cells, DA caused a concentration-dependent attenuation of forskolin-stimulated adenylate cyclase activity. Maximal inhibition was 27% of total activity, with an EC_{50} value of 624 nM (Fig. 10A; Table 1). Quinpirole was approximately as efficacious as dopamine; that is, the maximal inhibition induced by quinpirole was similar to that induced by DA. LY181990 was less potent and less efficacious than its active isomer (Fig. 10B; Table 1), indicating that D-2 receptor-mediated inhibition of adenylate cyclase activity was stereoselective. The finding that LY181990 caused measurable inhibition of enzyme activity in only 2 out of 3 experiments (Table 1) suggests that the compound has little or no agonist activity. Similarly, (-)3-PPP did not have detectable agonist activity. Bromocriptine, with an EC_{50} value of 45 nM, was the most potent agonist. Bromocriptine and (+)3-PPP were less efficacious than DA; thus, the drugs appeared to be partial agonists.

Inhibition of adenylate cyclase activity in LZR1 cells by $10 \mu\text{M}$ DA was prevented by including $10 \mu\text{M}$ (+)-butaclamol in the assay (Fig. 11), indicating that inhibition of DA is receptor-mediated. Also, treatment of LZR1 cells with pertussis toxin (50 ng/ml of growth medium for 16 hours) blocked DA-inhibited enzyme activity in membranes prepared from the cells (Fig. 10C), suggesting that G_i mediates inhibition of enzyme activity by DA. Interestingly, forskolin-stimulated adenylate cyclase activity in membranes from pertussis toxin-

treated cells was approximately 2.5-fold greater than activity in control membranes.

TABLE 1

**Inhibition of radioligand binding
and adenylate cyclase activity by agonists**

The apparent affinity (K_i) of 6 drugs for D-2 receptors on membranes prepared from LZR1 cells, determined by inhibition of the binding of [3 H]spiroperidol (0.2 nM), is shown, as well as the concentration of each drug that caused half-maximal inhibition of forskolin-stimulated adenylate cyclase activity (EC_{50}). Values for drug concentrations, expressed as μ M, are the geometric means of results from 3 experiments (EC_{50} , quinpirole and (+)3-PPP), 4 experiments (EC_{50} -DA) or 2 (K_i , all drugs; EC_{50} , bromocriptine) experiments. The maximal inhibition of adenylate cyclase activity observed (Max) is expressed as the mean \pm SEM of the percent inhibition of total activity in the presence of 10 μ M forskolin. For LY181990, the results shown are from two experiments in which inhibition of enzyme was observed. There was no inhibition in a third experiment.

<u>Drug</u>	<u>K_i</u>	<u>EC_{50}</u>	<u>Max</u>
Dopamine	17	0.6	27 \pm 3%
Quinpirole	9	0.7	28 \pm 2%
LY181990	277	5.0	10 \pm 2%
Bromocriptine	0.0024	0.04	17 \pm 1%
(+)3-PPP	33	4.0	16 \pm 3%
(-)3-PPP	0.87	--	--

Discussion

As reported previously, Ltk⁻ cells transfected with a rat D-2 receptor cDNA express a high density of DA D-2 receptors (1). We have characterized one subclone of these cells, designated LZR1, that stably expresses D-2

receptors at a density of 750 to 1000 fmol/mg of protein (present results; ref. 1).

As the D-2 cDNA is contained in the plasmid pZem3, under the regulation of a zinc-inducible promoter, the effect of zinc treatment on the properties of radioligand binding was determined. D-2 receptors on LZR1 cells appear to be maximally expressed in the absence of zinc, so that zinc treatment caused no increase in the density of receptors. The lack of responsiveness to zinc is not a characteristic of the Ltk⁻ cell line, since we have isolated a second transfected line of Ltk⁻ cells, LZR2, in which the density of D-2 receptors is elevated approximately 50% by treatment with zinc, from 435 to 630 fmol/mg of protein. The affinity of D-2 receptors for [³H]spiroperidol was not significantly altered by zinc treatment.

The apparent affinity of D-2 receptors on LZR1 cells for several agonists and related drugs was determined by inhibiting the specific binding of [³H]spiroperidol with the drugs in the presence of GTP. Bromocriptine was an extremely potent agonist, with a K_i value of 2 nM. The potency of quinpirole, like that of DA, was approximately 10 μM. The binding of agonists was stereoselective, since LY181990 was much less potent than its active enantiomer, quinpirole, and (-)-3-PPP was more potent than (+)-3-PPP. These data extend our previous investigation of the binding of antagonists to D-2 receptors on LZR1 cells (1).

Physiological effects of stimulation of D-2 receptors appear to be mediated by a guanine nucleotide-binding protein, G_i, that inhibits adenylate cyclase activity (16). High affinity binding of agonists is thought to represent a ternary complex composed of agonist, receptor, and the α-subunit of G_i (G_{ia}), and inhibition of agonist binding to D-2 receptors by GTP represents GTP-induced uncoupling of D-2 receptors from G_{ia}. To evaluate the ability of D-2 receptors encoded by

the RGB-2 cDNA to couple to G proteins, the effect of GTP on the potency of DA for inhibition of radioligand binding was determined. In preliminary studies using LZR1 cells, we found that in our standard assay buffer containing 120 mM NaCl and no added Mg^{2+} , the binding of DA was not sensitive to GTP (1), although under the same conditions the binding of DA to rat striatal membranes is inhibited by GTP (data not shown). There were three possible explanations for the lack of sensitivity to GTP in LZR1 membranes: (1) LZR1 cells, derived from Ltk⁻ cells, could lack the appropriate G-protein. (2) The RGB-2 cDNA could encode only a binding subunit of the D-2 receptor. This possibility seemed unlikely because of the similarity between the predicted primary structure of the protein encoded by RGB-2 and other receptors coupled to guanine nucleotide-binding proteins. (3) It could be that the ionic conditions of the binding assay were not appropriate for formation or destabilization of the ternary complex. In the studies described here, ionic conditions were varied to increase the likelihood of observing GTP-sensitive binding. To maximize high-affinity agonist binding, tissue was preincubated with $MgCl_2$, and $MgCl_2$ was included in the assay buffer (17). To maximize GTP-induced destabilization of the ternary complex, NaCl was added together with GTP (17). Under these conditions, addition of GTP and NaCl decreased the potency of DA for D-2 receptors and increased the slope of the inhibition curve in membranes from LZR1 cells. It is interesting that the ionic requirements for formation and destabilization of the ternary complex seem to be more stringent in membranes from LZR cells than in membranes from rat striatum.

Inhibition of adenylate cyclase activity by DA D-2 receptors is a well-characterized phenomenon (3, 18, 19). Inhibition of adenylate cyclase activity by several drugs was assessed in membranes from LZR1 cells. Maximally effective concentrations of DA and the D-2-selective

agonist quinpirole decreased forskolin-stimulated enzyme activity by almost 30%. Inhibition of enzyme activity by DA was blocked by the D-2 antagonist (+)-butaclamol. The most potent agonist tested, bromocriptine, appeared to be

5

a partial agonist, as reported by others (19, 20). Inhibition of adenylate cyclase by agonists was stereoselective, since LY181990, the dextrorotatory enantiomer of quinpirole, had little or no efficacy. As has been reported previously, the partial agonist (+)3-PPP is a stronger agonist than (-)3-PPP, although (-)3-PPP binds to D-2 receptors with higher affinity (21, 22).

10

We observed no inhibition of adenylate cyclase activity by (-)3-PPP in membranes from LZR1 cells. The EC_{50} values determined for inhibition of adenylate cyclase activity

15

by agonists were generally lower than the K_i values determined in assays of ligand binding (Table 1). This could be due to the presence of a receptor reserve on these cells, although the observation that drugs differed in maximal inhibition of enzyme activity suggests that there is not a large receptor reserve even for the most efficacious agonists, DA and quinpirole. Also, there would be no receptor reserve for a partial agonist such as (+)3-PPP. An alternative explanation is that

20

inhibition of adenylate cyclase activity could be related to the binding of agonists to D-2 receptors in a high-affinity state induced by formation of the ternary complex, as had been proposed for inhibition of enzyme activity by DA in the anterior pituitary (23). On the

25

other hand, K_i values in the present report, determined in the presence of GTP and NaCl, reflect the binding of agonists to receptors in a state of low affinity. Direct

30

comparisons are difficult, since adenylate cyclase and radioligand binding assays were carried out at different temperatures, but the K_i value for DA binding to the high-affinity class of sites ($0.3 \mu M$) is close to the EC_{50} value for DA-inhibited adenylate cyclase ($0.6 \mu M$),

35

whereas the K_i values for the binding of DA to the low-

affinity class of sites ($24 \mu\text{M}$) and binding in the presence of GTP ($17 \mu\text{M}$) are considerably higher.

DA did not inhibit adenylate cyclase activity in membranes from LZR1 cells that had been treated with pertussis toxin. Since pertussis toxin-catalyzed ADP-ribosylation of G_i prevents G_i -mediated inhibition of adenylate cyclase, this finding is consistent with the hypothesis that D-2 receptors interact with $G_{i\alpha}$ in the transfected LZR1 cells. As has been observed for stimulation of adenylate cyclase activity by isoproterenol after pertussis toxin-treatment of other cell types (24), treatment of intact LZR1 cells with pertussis toxin potentiated the ability of forskolin to stimulate adenylate cyclase activity, suggesting that in some cell lines G_i normally acts to attenuate forskolin- and hormone-stimulated adenylate cyclase activity.

We have characterized a cell line, transfected with the RGB-2 cDNA, that stably expresses a high density of D-2 receptors. With this cell line, it was determined the cDNA encodes a DA D-2 receptor that interacts productively with a guanine nucleotide-binding protein to inhibit adenylate cyclase activity. It seems likely that the RGB-2 cDNA would direct the expression of a functional D-2 receptor in almost any type of cell. For example, GH_4C_1 cells, derived from a rat pituitary tumor (25), are prolactin-secreting cells that lack DA receptors, even though lactotrophs in the rat anterior pituitary express D-2 receptors. Transfection of the RGB-2 cDNA into GH_4C_1 cells results in the expression of a D-2 receptor with functional characteristics similar to those described here (26). Cell lines created by transfection with a D-2 receptor cDNA will be useful in the study of mechanisms of action and regulation of D-2 receptors.

References for Example 1

1. Bunzow, J.R., H.H.M. Van Tol, D.K. Grandy, P. Albert, J. Salon, M. Christie, C.A. Machida, K.A. Neve, and O. Civelli. Cloning and expressing of a rat
5 D₂ dopamine receptor cDNA. Nature (Lond.)
336:783-787 (1988).
2. Kebabian, J.W., and D.B. Calne. Multiple receptors for dopamine. Nature (Lond.) 277:93-96 (1979).
3. De Camilli, P., D. Macconi, and A. Spada. Dopamine
10 inhibits adenylate cyclase in human prolactin-secreting pituitary adenomas. Nature (Lond.)
278:252-254 (1979).
4. Malgaroli, A., L. Vallar, F.R. Elahi, T. Pozzan, A. Spada, and J. Meldolesi. Dopamine inhibits
15 cytosolic Ca²⁺-increases in rat lactotroph cells: Evidence of a dual mechanism of action. J. Biol. Chem. 262:13920-13927 (1987).
5. Drouva, S.V., E. Rerat, C. Bihoreau, E. Laplante, R. Rasolonjanahary, H. Clauser, and C. Kordon.
20 Dihydropyridine-sensitive calcium channel activity related to prolactin, growth hormone, and luteinizing hormone release from anterior pituitary cells in culture. Interactions with somatostatin, dopamine, and estrogens.
25 Endocrinology 123:2762-2773 (1988).
6. Lacey, M.G., N.B. Mercuri, and R.A. North. Dopamine acts on D₂ receptors to increase potassium conductance in neurones of the rat substantia nigra zona compacta. J. Physiol. (Lond.) 392:397-
30 416 (1987).

7. Simmonds, S.H., and P.G. Strange. Inhibition of inositol phospholipid breakdown by D₂ dopamine receptors in dissociated bovine anterior pituitary cells. Neurosci. Lett. 60:267-272 (1985).
8. Enjalbert, A., F. Sladaczek, G. Guillon, P. Bertrand, C. Shu, J. Epelbaum, A. Garcia-Sainz, S. Jard, C. Lombard, C. Kordon, and J. Bockaert. Angiotensin II and dopamine modulate both cAMP and inositol phosphate productions in anterior pituitary cells: Involvement in prolactin secretion. J. Biol. Chem. 261:4071-4075 (1986).
9. Judd, A.M., I.S. Login, K. Kovacs, P.C. Ross, B.L. Spangelo, W.D. Jarvis, and R.M. MacLeod. Characterization of the MMQ cell, a prolactin-secreting cloned cell line that is responsive to dopamine. Endocrinology 123:2341-2350 (1988).
10. Lefkowitz, R.J. and M.G. Caron. Adrenergic receptors: Models for the study of receptors coupled to guanine nucleotide regulatory proteins. J. Biol. Chem. 263:4993-4996 (1988).
11. Uhler, M.D., and G.S. McKnight. Expression of cDNAs for two isoforms of the catalytic subunit of cAMP-dependent protein kinase. J. Biol. Chem. 262:15202-15207, 1987.
12. Gorman, C., R. Padmanabhan, and B.H. Howard. High efficiency DNA-mediated transformation of primate cells. Science 231:551-553 (1983).
13. Cheng, Y.-C. and W.H. Prusoff. Relationship between the inhibition constant (K_i) and the concentration of inhibitor which causes 50 per cent inhibition (I_{50}) of an enzymatic reaction. Biochem. Pharmacol. 22:3099-3108 (1973).
14. Salomon, Y., C. Londos, and M. Rodbell. A highly sensitive adenylate cyclase assay. Analyt. Biochem. 58:541-548 (1974).

15. Peterson, G.L. A simplification of the protein assay method of Lowry et al. which is more generally applicable. Analyt. Biochem. 83:346-356 (1977).
- 5 16. Cote, T.E., E.A. Frey, C.W. Grewe, and J.W. Kebebian. Evidence that the dopamine receptor in the intermediate lobe of the rat pituitary gland is associated with an inhibitory guanyl nucleotide component. J. Neural. Trans. Suppl. 10 18:139-147 (1983).
17. Hamblin, M.W., and I. Creese. ³H-Dopamine binding to rat striatal D-2 and D-3 sites: Enhancement by magnesium and inhibition by guanine nucleotides and sodium. Life Sci. 30:1587-1595 (1982).
- 15 18. Weiss, S., M. Sebben, J.A. Garcia-Sainz, and J. Bockaert. D₂-Dopamine receptor-mediated inhibition of cyclic AMP formation in striatal neurons in primary culture. Mol. Pharmacol. 27:595-599 (1985).
- 20 19. Onali, P., M.C. Olanas, and G.L. Gessa. Characterization of dopamine receptors mediating inhibition of adenylate cyclase activity in rat striatum. Mol. Pharmacol. 28:138-145 (1985).
20. Agui, T., N. Amlaiky, M.G. Caron, and J.W. Kebebian. Binding of [¹²⁵I]-N-(p-aminophenethyl)spiroperidol to the D-2 dopamine receptor in the neurointermediate lobe of the rat pituitary gland: A thermodynamic study. Mol. Pharmacol. 25 32:163-169 (1988).
- 30 21. Koch, S.W., B.K. Koe, and N.G. Bacopoulos. Differential effects of the enantiomers of 3-(3-hydroxyphenyl)-N-n-propylpiperidine (3-PPP) at dopamine receptor sites. Eur. J. Pharmacol. 92:279-283 (1983).
- 35 22. Meller, E., K. Bohmaker, Y. Namba, A.J. Friedhoff, and M. Goldstein. Relationship between receptor occupancy and response at striatal dopamine

- autoreceptors. Mol. Pharmacol. 31:592-598 (1987).
23. Borgundvaag, V., and S.R. George. Dopamine inhibition of anterior pituitary adenylate cyclase is mediated through the high-affinity state of the D_2 receptor. Life Sci. 37:379-386 (1985).
24. Abramson, S.N., M.W. Martin, A.R. Hughes, T.K. Harden, K.A. Neve, D.A. Barrett, and P.B. Molinoff. Interaction of β -adrenergic receptors with the inhibitory guanine nucleotide-binding protein of adenylate cyclase in membranes prepared from cyc-S49 lymphoma cells. Biochem. Pharmacol. 37:4289-4297 (1988).
25. Tashjian, A.H. Clonal strains of hormone-producing pituitary cells. Meth. Enzymol. 58:527-535 (1979).
26. Albert, P.R., K. Neve, J. Bunzow, and O. Civelli. Biological functions of the rat dopamine D_2 receptor cDNA expressed in GH_4C_1 rat pituitary cells. Proc. Endocrine Soc. 71:(in press).

Example 2

Summary

We have previously described a cDNA which encodes a binding site with the pharmacology of the D₂-dopamine receptor (Bunzow, J.R., et al. (19868) Nature 336, 783-787). We demonstrate here that this protein is a functional receptor, i.e., it couples to G-proteins to inhibit cAMP generation and hormone secretion. The cDNA was expressed in GH₄C₁ cells, a rat somatomammotrophic cell strain which lacks dopamine receptors. Stable transfectants were isolated and one clone, GH₄ZR₇, which had the highest levels of D₂-dopamine receptor mRNA on Northern blot, was studied in detail. Binding of D₂-dopamine antagonist ³H-spiperone to membranes isolated from GH₄ZR₇ cells was saturable, with K_d=96 pM, and B_{max}=2300 fmol/mg protein. Addition of GTP/NaCl increased the IC₅₀ value for dopamine competition for ³H-spiperone binding by two-fold, indicating that the D₂-dopamine receptor interacts with one or more G proteins. To assess the function of the dopamine binding site, acute biological actions of dopamine were characterized in GH₄ZR₇ cells. Dopamine decreased resting intra- and extracellular cAMP levels by 50-70% (EC₅₀=8±2 nM), and blocked completely VIP-induced enhancement of cAMP levels (EC₅₀=6±1 nM), which ranged from 8-12 times basal levels. Antagonism of dopamine-induced inhibition of VIP-enhanced cAMP levels by spiperone, (+)-butaclamol, (-)-sulpiride and SCH23390 occurred at concentrations expected from K_i values for these antagonists at the D₂-receptor and was stereo-selective. Dopamine (as well as several D₂-selective agonists) inhibited forskolin-stimulated

adenylate cyclase activity by $45 \pm 6\%$, with EC_{50} of 500-800 nM in GH₄ZR₇ membranes. Dopaminergic inhibition of cellular cAMP levels and of adenylylase activity in membrane preparations was abolished by pretreatment with pertussis toxin (50 ng/ml, 16h). Dopamine (200 nM) abolished VIP and TRH-induced acute prolactin release. These data show conclusively that the cDNA clone encodes a functional dopamine-D₂ receptor which couples to G proteins to inhibit adenylylase, and both cAMP-dependent and cAMP-independent hormone secretion. The GH₄ZR₇ cells will prove useful in elucidating further the biochemistry of the dopamine D₂ receptor.

Introduction

The major element controlling PRL¹ secretion from the pituitary is the concentration of dopamine in the hypophyseal portal bloodstream (1). Dopamine acts via dopamine-D₂ receptors on pituitary lactotrophs to inhibit basal and hormone-stimulated secretion of PRL (1-5). The dopamine-D₂ receptor interacts with pertussis toxin-sensitive, inhibitory G proteins (6-9) to reduce adenylylase activity, and to block enhancement of cAMP levels by other agents (6, 10-12). Dopamine also decreases $[Ca^{++}]_i$ in lactotrophs, and partially inhibits elevation of $[Ca^{++}]_i$ by other agents, such as TRH (13-15). Both dopaminergic inhibition of cAMP and of $[Ca^{++}]_i$ are mediated through coupling to one or more pertussis toxin-sensitive G proteins, and appear to contribute to dopamine inhibition of PRL secretion (15). The precise relation between these components of dopamine action has been difficult to study (15, 16) due to the presence of heterogeneous cell types, limitations of cell number, and variations in responsiveness of diverse lactotroph preparations.

The recent cloning of the dopamine-D₂ receptor cDNA (17) provides a useful tool to examine the intracellular actions and regulation of the receptor. To examine

whether the D₂-receptor clone directs synthesis of a functional receptor, and to define the pathway between dopamine-D₂ receptor activation and biological effect, we have transfected the dopamine-D₂ receptor cDNA into a pituitary-derived cell strain, GH₄C₁ cells. GH₄C₁ cells are rat pituitary cells which synthesize and secrete PRL and GH, and possess a variety of hormone, growth factor, and neurotransmitter receptors, second messenger systems, and ion channels and have provided an accessible model of lactotroph function (18, 19). However, these cells lack dopamine-D₂ receptors, which are present on normal lactotrophs, and thus provide an ideal host for studying the function of the dopamine-D₂ receptor. This report demonstrates that the gene product of the cDNA clone functions as a dopamine-D₂ receptor and couples to inhibitory G proteins to decrease cAMP accumulation² and PRL release. The GH₄C₁ transfectants characterized herein should provide a useful cell system in which the mechanisms of dopamine action at D₂ receptors may be studied further.

Experimental Procedures

Materials: Dopamine agonists and antagonists were from Research Biochemicals Incorporated (Waltham, MA), except quinpirole (Lilly), bromocryptine (Sandoz Research Institute), and (+) and (-)3-PPP (Astra). Rabbit antibody (lot CA-3) to 2-O-succinyl-cAMP-bovine serum albumin conjugate was obtained from ICN (Irvine, CA), rPRL standard and anti-rPRL antibody were from Dr. Salvatore Raiti, NIDDK, Bethesda, MD. Peptides were from Peninsula (, CA) or Sigma (St. Louis, MO). $\alpha^{32}\text{P}$ -dCTP (2,200 Ci/mmol), ^{125}I -2-O-(iodotyrosyl methyl ester)-succinyl cAMP (2,200 Ci/mmol), ^{125}I -rPRL (2,200 Ci/mmol), $\alpha^{32}\text{P}$ -ATP (10-50 Ci/mmol), ^3H -cAMP (31.9 Ci/mmol) were from New England Nuclear (Boston, MA). All other chemicals were reagent grade, obtained primarily from Sigma.

Methods:

Construction of pZEM-D₂-cDNA: The pZEM-3 plasmid (20) was cut at the *Bgl* II site between the metallothionein promotor and hGH 3'-flanking sequence. Full-length dopamine-D₂ cDNA (17) was excised from λ GT10 with *Sa*I and was ligated to the cut pZEM-3 plasmid in the presence of dATP and dGTP (250 μ M) and transformed into E. Coli strain XL-1 (Stratagene). Recombinants were characterized by their hybridization to ³²P-labelled D₂-cDNA, followed by restriction analysis and DNA sequencing. Recombinants with cDNA inserts in the sense orientation were prepared and purified by CsCl gradient centrifugation for transfection into eucaryotic cells.

Cell Culture: GH₄C₁ cells, obtained from Dr. A.H. Tashjian, Jr. (Harvard University, Boston, MA) and subclones were grown in Ham's F10 medium, supplemented with 10% fetal bovine serum, at 37°C in 5% CO₂. For studies of ³H-spiperone binding or adenylate cyclase activity, cells were grown in Dulbecco's modified Eagle's medium supplemented with 10% fetal bovine serum at 37°C in 10% CO₂. Media were changed 12-24h prior to transfection or experimentation. For transfection, GH₄C₁ cells were grown to 2-4 x 10⁶ cells/10 cm dish. 20 μ g of pZEM-D₂cDNA and 1 μ g pRSV-neo were co-precipitated with calcium phosphate in 2 ml of HEPES-buffered saline, and placed over cells for 10-20 min. 8 ml of warm F10 + 10% fetal calf serum (pH 7.0) was added, and the cells incubated for 4-5 h, 37°C. The medium was removed and 15% glycerol in HEPES-buffered saline was added, and incubated for 3 min., 37°C. The plates were rinsed and fresh F10 + 10% fetal calf serum added, and the cells placed in the incubator for 16-20 h. Fresh medium supplemented with 700 μ g/ml G418 (Geneticin, GIBCO, N.Y.) was added over the next 3-4 weeks, to select for stable transfectants expressing neo-resistance. Single colonies were isolated using sterile micropipette tips to take up individual colonies in 3-5 μ l. Once stocks of the

transfectant cell lines were stored frozen in liquid nitrogen, G418 was omitted from growth media.

RNA Isolation and Northern Blot Analysis: Cells were rinsed in calcium-free Hepes-buffered saline + 0.02% EDTA and extracted with Tris-buffered guanidinium hydrochloride, centrifuged (33,000 rpm, 16h) through a 1.7 g/ml CsCl pad, and the pellets extracted with phenol/chloroform and ethanol precipitated (21). RNA was resuspended and quantitated by UV absorbance at OD=260 nm. For Northern blots, RNA was denatured in glyoxal/dimethylsulfoxide (1h, 50°C), and run on a 1% agarose gel in 10 mM sodium phosphate. RNA was blotted overnight onto nylon membrane (N-bond, Amersham), baked at 80°C for 2h. Prehybridization was as described, for 6h at 42°C. Random-primed ³²P-labelled 1.6kb *Bam*HI-*Bgl*III fragment of the D₂-cDNA (1-2 x 10⁶ dpm/μg) was used for hybridization, 16-20h at 42°C in 50% formamide. Blots were washed in 2 x SSC for 10 min, room temperature, followed by 3 x 15 min. wash in 0.2 x SSC, 0.5% SDS, 70°C, and exposed to X-ray film overnight at -80°C, with intensifying screen.

Ligand Binding: Cell membranes were prepared by first replacing growth medium with ice-cold hypotonic buffer (1 mM Hepes, pH 7.4, 2 mM EDTA). After swelling for 10-15 min, the cells were scraped from the plate and centrifuged at 24,000g for 20 min, lysed with a Brinkman Polytron homogenizer at setting 6 for 10 sec in Tris-isosaline (50 mM Tris, pH 7.4, 0.9% NaCl) and stored at -70°C for receptor binding experiments, or resuspended in 50 mM Tris, pH 7.4, centrifuged as above, and resuspended in 50 mM Tris for immediate use in adenylate cyclase assay (below). For binding assays, the membrane preparation was thawed, centrifuged (24,000g x 20 min) and resuspended in Tris-isosaline except where indicated. Aliquots of membrane preparation were added to tubes containing 50 mM Tris, pH 7.4, 0.9% NaCl, 0.025% ascorbic acid, 0.001% bovine serum albumin, ³H-spiperone and

indicated drugs. (+)-Butaclamol ($2 \mu\text{M}$) was used to define nonspecific binding, which was less than 10% of total binding at concentrations of radioligand near the K_D value. Assays were carried out in duplicate, in a volume of 2 ml for saturation analyses or 1 ml for inhibition analyses. Incubations were initiated by addition of 10-50 μg of membrane protein, carried out at 37°C for 50 min, and stopped by addition of 10 ml of ice-cold buffer (10 mM Tris, pH 7.4, 0.9% NaCl) to each tube. The samples were immediately filtered through glass-fibre filters (Schleicher and Schuell No. 30) and washed with 10 ml of ice-cold buffer. Radioactivity retained on the filter was counted using a Beckman LS 1701 scintillation counter. In experiments to examine the effect of GTP on dopamine binding, cells were harvested, centrifuged, resuspended in Tris- Mg^{++} (50 mM Tris, pH 7.4, 4 mM MgCl_2) and incubated for 15 min, 37°C . After centrifugation, the resuspended protein was added to assays containing Tris- Mg^{++} with no added GTP or NaCl, or Tris- Mg^{++} with 120 mM NaCl and 100 μM GTP.

CAMP and PRL Assay: Cells were plated in 6-well, 35mm dishes, 3-7 days prior to experimentation. Cells were pre-incubated in 2 ml/well warm F10 + 0.1% (-)-ascorbic acid + 20 mM Tris (pH 7.2) (FAT) for 5-10 min, followed by addition of 1 ml/well of FAT + 100 μM IBMX + experimental compounds, and incubated for 30 min. at 37°C . Experimental compounds were diluted 200- to 1000-fold from stock solutions made immediately prior to assay. The final ethanol concentration never exceeded 0.1%, a concentration without effect on basal or VIP-enhanced cAMP or PRL levels in GH_4ZR_7 cells. Media were collected, and the cells were lysed immediately in 1 ml of boiling water. Cell lysates and media were centrifuged ($2000 \times g$, 10 min., 4°C), and the supernatants collected for assay as cell extracts. Cell extracts and media samples were frozen at -20°C until assay, if not assayed immediately. cAMP was assayed by a

specific radioimmunoassay as described (22), with antibody used at 1:500 dilution. After 16h incubation at 4°C, 20 µl of 10% BSA and 1 ml of 95% ethanol were added consecutively to precipitate the antibody-antigen complex. Standard curves showed IC₅₀ of 0.5±0.2 pmol using cAMP as standard. PRL was assayed in the media samples obtained as described above, except that IBMX was omitted during the 30 min incubation in FAT medium. PRL levels were determined by specific radioimmunoassay using Staphylococcus A lysate (IgSorb, The Enzyme Center, Malden, MA) to precipitate antigen-antibody complexes (23). Standard curves gave IC₅₀ of 36±6 ng using rPRL standard.

Adenylate Cyclase Assay: The conversion of α³²P-ATP to ³²P-cAMP was determined essentially as described by Salomon et al. (24). Membranes (10–50 µg) were added in a volume of 100 µl to an assay of 200 µl containing 50 mM Tris, pH 7.4, 5 mM cAMP, 1 mM IBMX, 1 mM MgCl₂, 0.5 mM EGTA, 0.25 mM ATP, 30 µM GTP, and about 2 × 10⁶ cpm of α³²P-ATP, and various drugs. Assays carried out in triplicate were initiated by warming to 25°C and terminated by cooling to 0°C. Trichloroacetic acid (100 µl of 30% solution) was added to each assay, and ³H-cAMP (30,000 cpm) was added to each tube as an internal standard. The assay volume was increased to 1 ml by addition of water, and tubes were centrifuged (2000g × 10 min). cAMP in the supernatant was isolated by sequential chromatography on columns containing Dowex AG50W-X4 resin and neutral alumina. The 2-ml eluate from each alumina column was dissolved in 10 ml of Bio-Safe II (RPI, Mount Prospect, IL) for liquid scintillation counting.

Calculations: Data from cAMP and PRL assays are expressed as means ± standard error for triplicate determinations. Curve-fitting parameters were obtained by nonlinear regression analysis using the Enzfitter program (Elsevier Biosoft). Average affinity, EC₅₀ and IC₅₀ values are geometric means of the indicated number of experiments. In competition experiments, K_i values were

calculated from experimentally determined IC_{50} values by the method of Cheng and Prusoff (25). All experiments were representative of 3-5 independent trials, with the exception of that presented in Figure 16 (2 trials).

5 Results

Characterization of Stable Transfectants: GH_4C_1 cells were cotransfected with pZEM- D_2 -cDNA and pRSV-neo, and colonies resistant to the antibiotic G418 were isolated and initially characterized by Northern blot analysis. One clone, GH_4ZR_7 , had higher levels of 2.5 kb D_2 mRNA than other clones (Figure 13A). Wild-type (untransfected) GH_4C_1 cells, as well as a GH_4C_1 cell transfectant (GH_4ZD_{10}) expressing the rat 5-HT $_{1A}$ receptor gene in the pZEM-3 vector³ showed no hybridization to the D_2 -receptor probe. Pretreatment of GH_4ZR_7 cells with 100 μM ZnSO $_4$ for 36h induced a marked enhancement of D_2 receptor mRNA, indicating that the transcribed mRNA is regulated by the zinc-sensitive metallothionein promoter (20). The GH_4ZR_7 clone was used for further analysis (below) because of the high levels of dopamine- D_2 receptor expression in this clone.

Specific binding of the selective dopamine- D_2 receptor antagonist, 3H -spiperone, was assayed in crude membranes prepared from GH_4ZR_7 cells (Figure 13B). The GH_4ZR_7 membranes showed a saturable component of 3H -spiperone binding which was displaced by 2 μM (+)-butaclamol, whereas membranes from wild-type GH_4C_1 cells showed no specific 3H -spiperone binding (data not shown). In 5 experiments, the GH_4ZR_7 cell membranes showed maximal specific 3H -spiperone binding of 2046 \pm 315 fmol/mg of protein, and a mean K_D value of 96 \pm 1 pM. These values demonstrate robust expression of dopamine- D_2 binding sites receptors in these cells with affinity for 3H -spiperone comparable to that obtained in rat striatal membranes, and in Ltk⁻ cells transfected with pZEM- D_2 -cDNA (17).

To ascertain whether the expressed dopamine binding site interacted with a G protein, inhibition of ^3H -spiperone binding by dopamine was assayed in GH_4ZR_7 cell membranes, in the absence or presence of 100 μM GTP and 120 mM NaCl (Figure 13C). Assays were carried out in the presence of 4 mM MgCl_2 to promote high affinity binding of dopamine. In the absence of added GTP and NaCl, dopamine inhibited ^3H -spiperone binding with $\text{IC}_{50}=49\pm15$ μM , and Hill coefficient of 0.69, suggesting the presence of high and low affinity sites for dopamine. Analyzing the data according to a model assuming the presence of two classes of binding sites indicated that 46 \pm 15% of the receptors had a high affinity ($K_D=0.5$ μM) for dopamine and the remaining receptors had lower affinity (30 μM) for the agonist. In the presence of GTP and NaCl, the IC_{50} for dopamine was shifted two-fold to 109 \pm 20 μM ($K_D=17$ μM) with Hill coefficient closer to unity (0.93). Thus, the presence of GTP/NaCl converts the dopamine receptors from a heterogeneous population of high and low affinity receptors to a nearly homogeneous population of receptors in a low-affinity agonist state, as observed in striatal membrane preparations (6, 7). These data suggest that the cloned dopamine binding site interacts with G proteins when expressed in GH_4 cells.

Dopamine Actions on cAMP and PRL Levels: To test directly the function of the expressed dopamine- D_2 receptor clone, the actions of dopamine on cellular cAMP levels were measured. These assays were conducted in the presence of 100 μM IBMX, to inhibit phosphodiesterase activity in these cells (22). Thus, the observed changes in cAMP levels reflect changes in the rate of synthesis of cAMP rather than changes in its degradation. Dopamine actions on basal cAMP levels were measured, as well as dopamine inhibition of VIP-enhanced levels of cAMP. GH_4C_1 cells respond to VIP with an enhancement of cAMP accumulation (Figure 14A) as described by others (22, 26). Dopamine had no effect on extracellular cAMP levels

in wild-type GH_4C_1 cells, whether VIP was omitted or present during the incubation. This result is consistent with the lack of D_2 -dopamine receptor mRNA and binding in GH_4C_1 cells (Figure 13), and indicates that these cells also lack a detectable D_2 -dopamine response since dopamine does not elevate cAMP concentrations. In media from GH_4ZR_7 cells, dopamine inhibited both basal cAMP levels (by 50-70%), and reduced VIP-enhanced cAMP to basal levels. These actions of dopamine to reduce cAMP levels were consistently observed in all experiments and dopamine was equally effective in lowering intracellular cAMP levels (Figure 14B). As observed previously in GH_4C_1 cells (22), both intra- and extracellular cAMP levels change in parallel, although changes in extracellular cAMP may be more pronounced due to lower recovery of extracted intracellular cAMP. Dopamine actions on cAMP accumulation were blocked by (-)-sulpiride, a highly selective dopamine- D_2 antagonist, whereas the inactive stereoisomer, (+)-sulpiride, did not block dopaminergic inhibition of cAMP accumulation. Stereo-selective blockade by sulpiride suggested that inhibition of cAMP levels in GH_4ZR_7 by dopamine was mediated by activation of a dopamine- D_2 receptor not present in wild-type GH_4C_1 cells.

The physiological outcome of dopamine action is inhibition of secretion, which was assayed by measuring acute (30 min) PRL release in GH_4ZR_7 cells (Figure 14D). VIP and TRH enhanced PRL secretion 1.5- and 3-fold, respectively. VIP is thought to enhance PRL release by a cAMP-dependent mechanism (22, 23, 26), while TRH acts by a cAMP-independent mechanism linked to calcium mobilization (19, 27). Dopamine did not inhibit basal PRL release, but both VIP- and TRH-induced enhancement of PRL secretion were blocked by dopamine. Thus, dopamine blocked both cAMP-dependent and cAMP-independent secretion in GH_4ZR_7 cells. These actions of dopamine were reversed by (-)-sulpiride, but not by (+)-sulpiride. In

GH₄C₁ cells, dopamine had no effect on basal, VIP-stimulated, or TRH-stimulated secretion of PRL (data not shown).

To examine whether concentrations required for biological response correlated with affinity for the dopamine-D₂ receptor, dose-response relations were examined for dopamine actions on cAMP levels (Figure 15). Dopamine potently inhibited intra- and extracellular levels of cAMP with similar EC₅₀ values. Furthermore, dopamine inhibited both basal and VIP-enhanced cAMP accumulation with EC₅₀ values of 8±2 nM and 6±1 nM, respectively. These data demonstrate that dopamine inhibits both basal and stimulated cAMP accumulation with approximately equal potency. The high potency of these inhibitory actions of dopamine supports the assertion that GH₄ZR₇ cells express a functional dopamine-D₂ receptor.

The pharmacological specificity of dopaminergic inhibition of VIP-enhanced levels of cAMP in GH₄ZR₇ was examined further using specific receptor antagonists (Figure 16). The data show that certain receptor antagonists reverse dopamine-induced inhibition of VIP-enhanced levels of extracellular cAMP. Maximal cAMP (100%) corresponded to cAMP levels in the presence of VIP alone. Low concentrations of dopamine-D₂ antagonists (spiperone, (+)-butaclamol, (-)-sulpiride) blocked dopamine action, whereas SCH23390, a specific dopamine-D₁ antagonist, was active only at very high concentrations. Inactive stereoisomers of D₂-antagonists ((-)-butaclamol, (+)-sulpiride (Figure 14C)) had little or no effect on dopamine action. Antagonists added in the absence of dopamine did not alter cAMP concentrations. Estimated K_i values obtained from IC₅₀ values for the antagonists (see description of Figure 16) were similar to values determined from binding competition studies of the dopamine D₂ receptor (17), showing that inhibition of cAMP levels by dopamine in

GH₄ZR₇ cells is mediated by a receptor which is pharmacologically indistinguishable from the dopamine-D₂ receptor.

Inhibition of Adenylate Cyclase: To assess directly inhibition of adenylate cyclase activity by dopamine receptor agonists, the conversion of ³²P-ATP to ³²P-cAMP was measured in membranes prepared from GH₄ZR₇ cells (Figure 17). Dopamine inhibited total forskolin (10 μM)-stimulated activity by 45% with an average EC₅₀ value of 0.36 μM (n=5). As observed in pituitary (11) and striatal (28) membranes, bromocryptine behaved as partial agonist, maximally inhibiting enzyme activity by 23% (EC₅₀=6 nM). Inhibition of adenylate cyclase activity by selective D₂-agonists was stereo-selective. For example, quinpirole inhibited forskolin-stimulated cyclase activity by 41% (EC₅₀=0.32 nM), whereas LY181990, the inactive (+)-enantiomer of quinpirole, cause no consistent reduction in enzyme activity. Similarly, (+)-3-PPP (EC₅₀=0.86 nM) was as efficacious as dopamine, whereas the enantiomer (-)-3-PPP did not consistently reduce adenylate cyclase activity. VIP also stimulated adenylate cyclase activity in GH₄ZR₇ cell membranes, as reported for wild-type GH₄C₁ cell membranes (29). Total activity stimulated by 200 nM VIP was 22±7 pmol/mg protein/min (n=3), and VIP-enhanced activity was inhibited 41% by dopamine (100 μM), compared to 50-55% inhibition in rat anterior pituitary membranes (11). No effect of dopamine on basal adenylate cyclase activity was observed in these preparations. Nevertheless, inhibition by dopamine of forskolin- or VIP-stimulated adenylate cyclase activity provides a likely mechanism for inhibition of cAMP accumulation by dopamine in GH₄ZR₇ cells.

Pertussis Toxin Sensitivity: Sensitivity to pertussis toxin is a hallmark of receptors, such as the dopamine-D₂ receptor (6-12, 15), which couple to inhibitor G proteins (e.g., G_i or G_o) to induce responses.

Pretreatment of GH₄ZR₇ cells with pertussis toxin for 16h (Fig. 12) uncoupled dopamine-mediated inhibition of forskolin-stimulated membrane adenylate cyclase activity, and abolished inhibition of basal and VIP-stimulated cAMP accumulation by dopamine. The concentration of pertussis toxin and incubation time used produce maximal blockage of somatostatin responses in wild-type cells (30), and the dopamine responses were almost completely inhibited under these conditions. By contrast, basal and VIP-stimulated cAMP accumulation, as well as basal and forskolin-stimulated cyclase activity, were not significantly altered by pertussis toxin pretreatment. These data support the assertion that the expressed cDNA clone codes for a dopamine-D₂ binding site which is functionally coupled to inhibitory G proteins present in GH₄ cells, and thus represents a bona fide receptor.

Discussion

The cDNA clone coding for a dopamine-D₂ binding site (17) was expressed in GH₄C₁ cells to determine whether the clone expresses a functional D₂ receptor, which is coupled by pertussis toxin-sensitive inhibitory G proteins to inhibition of adenylate cyclase, cAMP accumulation, and inhibition of PRL secretion (6-12, 15). Dopamine-D₂ receptors were expressed specifically from a D₂-cDNA construct under the regulation of the mouse metallothionein promoter (20), as evidenced by the presence of D₂ receptor mRNA in the GH₄ZR₇ transfectant. Dopamine-D₂ receptor mRNA levels were undetectable in untransfected GH₄C₁ cells, or in cells transfected with the rat 5-HT_{1A} receptor subtype (Figure 13A). The mRNA species found in GH₄ZR₇ cells was approximately the same molecular weight as D₂ receptor mRNA found in rat brain (17). Levels of dopamine-D₂ mRNA increased by addition of 100 μ M Zn⁺⁺ in GH₄ZR₇ cells, indicating that expression of the mRNA is controlled by the Zn⁺⁺-sensitive mouse metallothionein promoter, and does not represent

transcription of the endogenous gene. Specific binding of ^3H -spiperone binding was present only in GH_4ZR_7 cells, and was increased by $100\ \mu\text{M}\ \text{Zn}^{++}$, and thus correlated with the expression of dopamine- D_2 receptor mRNA in these cells. The D_2 -receptor mRNA was transcribed to yield robust expression of high affinity dopamine- D_2 binding sites in GH_4ZR_7 cells.

The presence of dopamine- D_2 binding in the GH_4ZR_7 transfectant correlated with potent and powerful inhibition of cAMP accumulation and PRL release, as well as inhibition of forskolin-stimulated adenylate cyclase activity, actions of dopamine not observed in untransfected GH_4C_1 cells. These inhibitory actions of dopamine match exactly the known physiological actions of dopamine in pituitary lactotrophs (1, 2). In particular, dopamine controls PRL secretion and cAMP accumulation in lactotrophs such that stimulation of these processes does not occur unless dopamine concentrations decrease to low levels (1, 3-5). Similarly, in the presence of maximal concentrations of dopamine, VIP does not enhance cAMP levels or PRL secretion in GH_4ZR_7 cells (Figure 14). The potency of dopamine inhibition of basal and VIP-enhanced cAMP accumulation in GH_4ZR_7 cells (Figure 15) was in the range of concentration expected for lactotrophs, given that dopamine concentrations in hypophyseal portal blood vary from 7 nM in female rats during proestrous, to 20 nM during estrous, and are 3 nM in male rats (31). Detailed analysis of the pharmacology of dopamine-induced inhibition of adenylate cyclase and cAMP accumulation using specific agonists and antagonists are fully consistent with the conclusion that dopaminergic actions in GH_4ZR_7 cells are mediated by a receptor indistinguishable from the dopamine- D_2 receptor.

The discrepancy between the measured affinity of dopamine (Figure 13C), and the potency of dopamine to inhibit cAMP accumulation (Figure 13) raises the possibility that GH_4ZR_7 cells have "spare" receptors,

i.e., a sufficient excess of binding sites to shift the EC_{50} for biological action to values lower than the K_D value. An alternative explanation is that the receptor in membrane preparations has a lower affinity for agonists (but unchanged affinity for antagonists since measured IC_{50} values correlated with K_i values) than in intact cells. Since cytosolic or membrane-associated components present in intact cells are not entirely replaced in membrane binding and adenylate cyclase assay conditions, it is possible that components which allow for optimal function of the dopamine receptor in membrane preparations are lacking. This assertion is supported by the EC_{50} value for inhibition of particulate adenylate cyclase by dopamine (360 nM), which is close to K_i values obtained for dopamine from binding competition experiments (500 nM), but 100-fold higher EC_{50} values (6-8 nM) obtained for inhibition of CAMP levels by dopamine in intact cells. This difference in conditions may explain the observed differences between assays in intact versus particulate preparations. However, affinities of antagonists correlated well with estimated K_i values obtained from CAMP accumulation experiments (Figure 16), indicating that antagonist binding is similar in membranes and whole cells.

Further evidence of coupling of the expressed dopamine- D_2 binding site to G proteins is the shift of dopamine binding affinity to lower affinity in the presence of GTP. Such GTP-induced shifts in affinity have been reported for dopamine binding in membranes from rat brain (6, 7), and are due to interaction of the receptor with G proteins (8, 9). In the presence of GTP, the G protein dissociates from the receptor leaving the receptor in a low affinity agonist state (6, 7). In the case of dopamine, the difference between the affinities of the two states is small, hence the GTP-induced shift to the low-affinity state is small (two-fold) and requires the presence of Na^+ ion to maximize dissociation

of the G protein (28). The observation of a GTP-induced shift in dopamine affinity suggests that the expressed dopamine receptor is associated with one or more G proteins in GH₄ZR₇ cell membranes.

5 Although the inhibitory actions of dopamine indicate that the receptor couples to inhibitory G proteins, this was tested more directly by pretreating GH₄ZR₇ cells with pertussis toxin to inactivate inhibitory G proteins (30). Pertussis toxin pretreatment completely blocked dopamine
10 actions, without altering basal or VIP-stimulated cAMP accumulation. Thus, pertussis toxin prevented the dopamine-enhanced transduction of biological responses, presumably by uncoupling the dopamine-D₂ receptor from inhibitory G proteins. Unlike other systems (32), but as
15 seen in wild-type GH₄C₁ cells (30), enhancement of cAMP levels by stimulators (e.g., VIP) was not augmented in pertussis toxin-treated GH₄ZR₇ cells, nor were basal cAMP levels altered by pertussis toxin. This suggests that
20 inhibitory G proteins present in GH₄ZR₇ cells do not inhibit cAMP generation tonically.

 The presence of somatostatin receptors on GH₄C₁ cells (33) allows for direct comparison of the inhibitory actions of somatostatin and dopamine. Like dopamine, somatostatin does not inhibit basal adenylate cyclase
25 activity (29) or basal PRL secretion (30). Somatostatin-induced inhibition of VIP-stimulated cyclase activity (29), VIP-enhanced cAMP accumulation (22), and PRL secretion (23, 30) in GH₄C₁ cells were all about half the maximal inhibition induced by dopamine in GH₄ZR₇ cells.
30 Indeed, in GH₄ZR₇ cells, somatostatin was half as effective as dopamine at inhibiting VIP-enhanced cAMP accumulation (data not shown). The larger effects of dopamine were not due to the presence of an excessive number of D₂ receptors since the receptor number under
35 conditions of cAMP accumulation and PRL secretion experiments was less than 10,000 sites/cell, smaller than somatostatin receptor number (13,000 sites/cell) in GH₄C₁

cells (33). It is apparent that the dopamine receptors expressed in these cells are more effective at transducing inhibitory actions than somatostatin receptors, suggesting a more effective coupling of the dopamine receptor to the G proteins present in GH cells.

A second reason for expressing the dopamine- D_2 receptor cDNA in GH_4C_1 cells was to establish a new model system in which to study dopamine actions and D_2 receptor mechanisms. Current studies on dopamine action in rat lactotrophs, the most accessible cell system expressing dopamine- D_2 receptors until recently (16), have come to divergent conclusions on mechanisms of dopamine inhibition. In neurons, dopamine hyperpolarizes membrane potential (34) by opening potassium channels (35), actions mediated by coupling of the D_2 receptor to inhibitory G proteins (15). Membrane hyperpolarization induced by dopamine in lactotrophs (36) would close calcium channels, decreasing basal calcium influx and explaining observed decreases in basal $[Ca^{++}]_i$ induced by dopamine (13-15, 37, 38). However, dopamine has been observed to increase $[Ca^{++}]_i$ in certain pituitary cells (37). By examining dopamine-induced changes in $[Ca^{++}]_i$ in GH_4ZR_7 cells it will be possible to specifically associate the D_2 receptor with changes in $[Ca^{++}]_i$, and to ascertain the role of $[Ca^{++}]_i$ in mediating dopamine inhibition of hormone secretion. Another unresolved issue is the mechanism by which dopamine inhibits enhancement of secretion and by calcium-mobilizing hormones such as TRH. While some report inhibition by dopamine of TRH-induced enhancement of phosphatidyl inositol turnover (39, 40) and $[Ca^{++}]_i$ (13, 14), others find no change (38, 41). The observed inhibition by dopamine of TRH-induced PRL release in GH_4ZR_7 cells (Figure 14C) suggests that dopamine may be coupled by G proteins to processes (e.g., opening of potassium channels) which alter TRH-induced calcium-mobilization or phosphatidyl inositol turnover. GH_4ZR_7 cells will provide a homogeneous, abundant, and

highly-responsive preparation in which to study these and other questions regarding mechanisms of the dopamine-D₂ receptor.

5 The data presented in this report indicate that the
expressed clone (17) possesses the pharmacology of the
dopamine-D₂ receptor in all actions investigated including
inhibition of PRL secretion, cAMP generation, and
adenylate cyclase activity. The D₂ clone meets five basic
10 criteria for classification as a functional G protein-
coupled receptor: 1) the cDNA clone for the dopamine-D₂
binding site possesses the archetypical structure of G
protein-coupled receptors; 2) the clone expresses a
protein with saturable and specific binding properties;
3) agonist binding affinity to the expressed binding site
15 is decreased in the presence of GTP; 4) the expressed
receptor is coupled to functions (e.g., inhibition of
adenylate cyclase) known to be regulated by G proteins;
5) agents (e.g., pertussis toxin) which uncouple G
protein function uncouple activation of the expressed
20 receptor from generation of the appropriate response. In
conclusion, the interaction of the cloned dopamine-D₂
receptor with G proteins is productive, leading to
activation of α_i or α_o subunits and consequent inhibition
of cAMP and PRL levels by both cAMP-dependent and cAMP-
25 independent mechanisms.

References for Example 2

1. Ben-Jonathan, N. (1985), Endocr. Rev. 6, 564-589.
2. Memo, M., Missale, C., Carruba, M.O., and Spane, P.F. (1986), J. Neural Transm. (Suppl.) 22, 19-32.
- 5 3. Martinez de la Escalera, G., and Weiner, R.I. (1986), Endocrinology 123, 1682-1687.
4. Lopez, F.J., Dominguez, J.R., Sanchez-Criado, J.E., and Negro-Vilar, A. (1988), Endocrinology 124, 527-535.
- 10 5. Ibid, pp. 536-542.
6. Sibley, D.R., DeLean, A., and Creese, I. (1982), J. Biol. Chem. 257, 6351-6361.
7. DeLean, A., Kilpatrick, B.F., and Caron, M.G. (1982), Mol. Pharmacol. 22, 290-297.
- 15 8. Senogles, S.E., Benovic, J.L., Amlaiky, N., Unson, C., Milligan, G., Vinitzky, R., Spiegel, A.M., Caron, M.G. (1987), J. Biol. Chem. 262, 4860-4867.
9. Ohara, K., Haga, K., Bernstein, G., Haga, T., Ichiyama, A., and Ohara, K. (1988), Mol. Pharmacol. 33, 290-296.
- 20 10. DeCamilli, P., Macconi, D., and Spada, A. (1979), Nature 278, 252-254.
11. Onali, P., Schwartz, J.P., and Costa, E. (1981), P.N.A.S. 78, 6531-6534.
- 25 12. Cronin, M.J., Myer, G.A., MacLeod, R.M., and Hewlett, E. (1983), Am. J. Physiol. 244, E499-E504.
13. Schofield, J.G. (1983), F.E.B.S. Lett. 159, 79-82.
14. Margaroli, A., Vallar, L., Elahi, F.R., Pozzan, T., Spada, A., and Meldolesi, J. (1987), J. Biol. Chem. 261, 13920-13927.
- 30

15. Vallar, L. and Meldolesi, J. (1989), TIPS 10, 74-77.
16. Judd, A.M., Login, I.S., Kovacs, K., Ross, P.C.,
Spangelo, B.L., Jarvis, W.D., and MacLeod, R.M.
(1988), Endocrinology 123, 2341-2350.
- 5 17. Bunzow, J.R., VanTol, H.H.M., Grandy, D.K., Albert,
P., Salon, J., Christie, M., Machida, C., Neva,
K.A., and Civelli, O. (1988), Nature 336, 783-
787.
- 10 18. Tashjian, A.H., Jr. (1979), Meth. Enzymol. 58, 526-
535.
19. Ozawa, S., and Sand, O. (1986), Physiol. Rev. 66,
887-952.
20. Uhler, M., and McKnight, G.S. (1987), J. Biol. Chem.
262, 15202-15207.
- 15 21. Chirgwin, J.M., Prybyla, A.E., MacDonald, R.J., and
Rutter, W.J. (1979).
22. Dorflinger, L.J., and Schonbrunn, A. (1983),
Endocrinology 113, 1541-1550.
- 20 23. Dorflinger, L.J., and Schonbrunn, A. (1983),
Endocrinology 113, 1551-1558.
24. Salomon, Y., Londos, C., and Rodbell, M. (1974),
Analyt. Biochem. 58, 541-548.
25. Cheng, Y.-C., and Prusoff, W.H. (1973), Biochem.
Pharmacol. 22, 3099-3108.
- 25 26. Gourdji, D., Bataille, D., Vauclin, N., Grouselle,
D., Rosselin, G., and Tixier-Vidal, A. (1979),
FEBS Lett. 104, 165-168.
27. Albert, P.R., and Tashjian, A.H., Jr. (1984), J.
Biol. Chem. 259, 15350-15363.
- 30 28. Onali, P., Olinas, M.C., and Gessa, G.L. (1985),
Mol. Pharmacol. 28, 138-145.
29. Koch, B.D., and Schonbrunn, A. (1984), Endocrinology
114, 1784-1790.
- 35 30. Koch, B.D., Dorflinger, L.D., and Schonbrunn, A.
(1985), J. Biol. Chem. 260, 13138-13145.

31. Ben-Jonathan, N., Oliver, C., Weiner, H.J., Mical, R.S., and Porter, J.C. (1977), *Endocrinology* 100, 452-458.
- 5 32. Kurose, H., Katada, T., Amano, T., and Ui, M. (1983), *J. Biol. Chem.* 258, 4870-4875.
33. Schonbrunn, A., and Tashjian, A.H., Jr. (1978), *J. Biol. Chem.* 253, 6473-6483.
34. Bunney, B.S., Aghajanian, G.K., and Roth, R.H. (1973), *Nature (New Biol.)* 245, 123-125.
- 10 35. Lacey, M.G., Mercuri, N.B., and North, R.A. (1987), *J. Physiol.* 392, 397-416.
36. Taraskevich, P.S., and Douglas, W.W. (1978), *Nature* 276, 832-834.
- 15 37. Winiger, B.P., Wuvarin, F., Zahan, G.R., Wollheim, C.B., and Schlegel, W. (1987), *Endocrinology* 121, 2222-2228.
38. Law, G.J., Pachter, J.A., and Dannies, P. (1988), *Mol. Endocrinol.* 2, 966-972.
- 20 39. Journot, L., Homburger, V., Pantaloni, C., Priam, M., Bockaert, J., and Enjalbert, A. (1987), *J. Biol. Chem.* 262, 15106-15110.
40. Vallar, L., Vicentini, L.M., and Meldolesi, J. (1988), *J. Biol. Chem.* 263, 10127-10134.
- 25 41. Canonico, P.L., Jarvis, W.D., Judd, A.M., and MacLeod, R.M. (1986), *J. Endocrinol.* 110, 389-393.

Footnotes

¹The abbreviations used are: PRL, prolactin; GH, growth hormone; $[Ca^{++}]_i$, cytosolic free calcium concentration; VIP, vasoactive intestinal peptide; TRH, thyrotropin-releasing hormone; IBMX, 3-isobutyl-1-methyl xanthine; K_d , equilibrium dissociation constant; EC_{50} (IC_{50}), concentration required to elicit a half-maximal effect (inhibition).

²These results have been presented in part at the 71st Annual Endocrine Meetings, Seattle, WA, Abstract #1278.

³Albert, P.R., et al., manuscript in preparation.

Example 3

Abstract

A clone encoding a human dopamine D₂ receptor was isolated from a pituitary cDNA library and sequenced. The deduced protein sequence is 96% identical with that of the cloned rat receptor [Bunzow et al. (1988) Nature 336, 783-787] with one major difference: The human receptor contains an additional 29 amino acids in its putative third cytoplasmic loop. Southern blotting demonstrated the presence of only one human dopamine D₂ receptor gene. Two overlapping phage containing the gene were isolated and characterized. DNA sequence analysis of these clones showed that the coding sequence is interrupted by six introns and that the additional amino acids present in the human pituitary receptor are encoded by a single exon of 87-basepairs. The involvement of this sequence in alternative splicing and its biological significance are discussed.

Introduction

Dopamine neurons in the vertebrate central nervous system are involved in the initiation and execution of movement, the maintenance of emotional stability, and the regulation of pituitary function. Several human neurological diseases, including Parkinson's Disease (1) and schizophrenia (2), are thought to be manifestations of imbalances between dopamine receptors and dopamine. The receptors which mediate dopamine's effects have been divided into D₁ and D₂ subtypes, which are distinguished by their G-protein coupling (3, 4), ligand specificities, anatomical distribution and physiological effects (5).

The dopamine D₂ receptors have been of particular clinical interest due to their regulation of prolactin secretion (6) and their affinity for antipsychotic drugs (7, 8).

5 The dopamine D₂ receptors belong to the family of G-protein coupled receptors. The sequence similarity shared by members of this family enabled us to clone a rat brain dopamine D₂ receptor cDNA (9). We have used that clone to isolate the human pituitary dopamine D₂ receptor cDNA described here. We have found that the deduced amino acid sequences of these two receptors are very similar, with one notable difference. The human receptor contains an additional 29 amino acids in its putative third cytoplasmic loop which are encoded by one of the gene's exons. This genomic organization suggests that the existence of two dopamine D₂ receptor mRNAs is the result of an alternative splicing event.

Materials and Methods

Cloning of the Human Pituitary cDNA

Human pituitary tissue was a generous gift from Drs. N. Seidah and M. Chretien, Clinical Research Institute of Montreal, Canada. Poly(A)⁺ mRNA and cDNA were prepared as previously described (9). The cDNA was size-selected (1-6 kb) on agarose gels, isolated using Geneclean (Bio 101), ligated to EcoRI adaptors, cloned into λGT10 arms (Stratagene), and packaged (Gigapak Gold). Recombinants (1.5 x 10⁶) were screened on replica nylon filters (DuPont Plaque/Colony Hybridization filters) with [³²P]-labelled hybridization probes. Prehybridization and hybridization were performed in 50% formamide, 1% SDS, 2 x SSC (1 x SSC = 0.15 M NaCl/0.015 M sodium citrate, pH 7) at 37°C. The complete sequence of both strands of DNA were determined in M13mp19 using Sequenase (U.S. Biochemical) primed with synthetic oligonucleotides.

Expression and Pharmacology

The 2.5-kb human pituitary cDNA (hPitD₂) was cloned into pZem3 (a gift from Dr. E. Mulvihill, Zymogenetics) and co-transfected with the pRSVneo gene into mouse Ltk⁻ cells by CaPO₄ precipitation (10). A stable transfectant (L-hPitD₂Zem) was selected and maintained in 750 µg/ml of G418 (Geneticin sulphate, Gibco). Twenty hours prior to the harvesting of membranes, these cells were incubated with 70 µM zinc sulphate. Membranes were prepared from L-HPitD₂Zem, from the Ltk⁻ cell line expressing the cloned rat dopamine D₂ receptor (L-RGB2Zem-1) and from freshly dissected rat striata (Taconic Farm, Germantown, New York), as previously described (11, 12). For the binding assays, membrane protein was used at 10-15 µg from L-hPitD₂Zem, 50-75 µg from L-RGB2Zem-1, and 220-250 µg from rat striatum. The binding assays were incubated at 37°C for 60 minutes in 50 mM Tris-HCl, 120 mM NaCl, 5 mM KCl, 2 mM CaCl₂, 1 mM MgCl₂, pH 7.4, and then stopped by rapid filtration over glass fiber filters (Schleicher and Schuell, No. 32) which had been presoaked in 0.5% polyethyleneimine. The filters were washed twice in ice cold 50 mM Tris-HCl, pH 7.4. Saturation curves were generated using increasing concentrations of the highly D₂-specific antagonist [³H]-domperidone (13). Antagonist drugs were evaluated for their ability to inhibit specifically bound [³H]-domperidone (1 nM). The B_{max}, K_d, and IC₅₀ values were determined as previously described (14).

Southern Blotting

Human genomic DNA prepared from a normal male donor was a gift from M. Litt. Three micrograms of DNA were digested with restriction enzymes and the fragments were electrophoresed in 0.7% agarose and blotted onto nitrocellulose filters (Schleicher and Schuell). Prehybridization and hybridization were performed at 37°C in 50% formamide as previously described (15).

Genomic Sequencing

Genomic bacteriophage lambda libraries, prepared from normal human male DNA, were purchased from Stratagene and Clontech Laboratories, Inc. and screened with portions of the cloned rat dopamine D₂ receptor cDNA. DNA sequence was determined by a genomic sequencing approach (16, 17). Briefly, for each restriction enzyme used, cloned genomic phage DNA (50 µg) was digested, subjected to chemical cleavage, and the resulting fragments resolved in a denaturing polyacrylamide gel. The DNA was then transferred from the gel and immobilized onto nylon filters (Plasco Genetran). Using [³²P] end-labelled synthetic oligomers, ladders of sequence were visualized within exons and read into neighboring introns. The filters were then either reprobbed with a different oligomer, or a new filter was made in order to read the complementary sequence back across the exon. Both strands of the coding region were sequenced.

Results

Cloning and Sequence Analysis of the Human Pituitary cDNA

Using the rat brain D₂ receptor cDNA as probe, three partial cDNAs were isolated from a human pituitary library and sequenced. Two oligonucleotide probes based on these sequences were used to isolate a fourth cDNA, hPitD₂, which encoded a full-length receptor protein (Fig. 18). The human pituitary receptor contains seven putative transmembrane domains and lacks a signal sequence. Overall, the human and rat nucleotide sequences are 90% similar and show 96% identity at the amino acid level. Several consensus sequences for N-linked glycosylation, protein kinase A phosphorylation and palmitoylation (18) are conserved between the human and rat receptors. There are also 18 amino acid differences (including one deletion) between these proteins, and, strikingly, the human pituitary receptor

contains an additional 29 amino acids in its putative third cytoplasmic loop.

Expression and Pharmacological Evaluation of hPit D₂ cDNA

In order to evaluate the pharmacological characteristics of the human pituitary receptor, its cDNA was subcloned into pZem3 and expressed in mouse Ltk⁻ cells (L-hPitD₂Zem). Membranes prepared from these cells showed specific binding of [³H]-domperidone, a D₂-selective antagonist (12, 13), with a B_{max} of 4.05 +/- 0.3 pmol per mg (n=2) protein and a K_d of 0.74 +/- 0.11 nM (n=2). This K_d value is in excellent agreement with the published value of 0.74 nM in mouse brain membranes (13). A Scatchard plot of the data was linear. There was no detectable [³H]-domperidone binding in membranes prepared from cells transfected with pZem3 alone (data not shown). [³H]-domperidone binding to L-hPitD₂Zem membranes was inhibited by a number of dopamine D₂-specific drugs (Fig. 19). Their rank order of potency was: spiperone, (+)-butaclamol, haloperidol, and sulpiride. The serotonin-selective antagonist mianserin and the D₁-selective antagonist SCH-23390 inhibited domperidone binding only at very high concentrations, as did the inactive isomer (-)-butaclamol. These values are essentially identical to those obtained with membranes from Ltk⁻ cells transfected with the cloned rat dopamine D₂ receptor cDNA (L-RGB2Zem-1) and from rat striatum (Fig. 22).

Human Dopamine D₂ Receptor Gene

Using portions of the rat cDNA as probe, a clone was isolated from a human genomic library. This genomic clone, λHD2G1, contained a 1.6-kb BamHI fragment which encoded the last 64 amino acids of the human D₂ receptor and 1.2-kb of 3' non-coding sequence. The 1.6-kb fragment was used to probe a Southern blot of human genomic DNA digested with three restriction enzymes.

Each enzyme generated a single fragment that hybridized to the probe (Fig. 20), indicating that there is probably only one human dopamine D₂ receptor gene.

In order to isolate a genomic clone that encoded the N-terminus of the human receptor protein, a 118-bp restriction fragment from the cloned rat dopamine D₂ receptor cDNA (corresponding to amino acid residues 1-39) was used to screen a second genomic library. λ HD2G2 was isolated and found to overlap with λ HD2G1 by 400 nucleotides (Fig. 21a). Together, these phage span 34-kb of the human dopamine D₂ receptor gene locus, DRD2 (19), and contain the sequence found in the hPitD₂ cDNA plus sequences that extend 15-kb downstream of the polyadenylation signal and 3.7-kb upstream of the translation initiation site. To characterize the intron/exon structure of the gene, a genomic sequencing approach employing oligonucleotide probes and chemical cleavage was used (Fig. 21b). Since the divergence of nucleotide sequences between human and rat members of this receptor family is approximately 10% (unpublished observations), we were able to initiate the genomic sequencing relying on hybridization probes and restriction sites that are present in the cloned rat dopamine D₂ receptor cDNA. Our results demonstrate that the coding portion of the human dopamine D₂ receptor gene is divided into seven exons (Fig. 21b). Interestingly, we found that exon five is 87-bp long and encodes the entire 29 amino acid sequence present in the cloned human pituitary receptor (Figs. 18 and 21c). Analysis of the six introns revealed that each contains acceptor and donor sequences that conform to the GT/AG rule (20), as summarized in Fig. 23. The approximate sizes of the introns were based on the results of Southern blotting experiments (data not shown). When compared, the genomic and cDNA sequences were found to differ by only two silent transitions, one at 939 (T to C in the gene) and the other at 957 (C to T in the gene).

Discussion

Several alternative hypotheses might account for the extra sequence present in our human pituitary D₂ receptor cDNA clone. One possibility is that the human gene contains the extra 87 bases and that the rat gene does not. Another is that both human and rat have two distinct genes which code for two different dopamine D₂ receptors. Finally, alternative splicing of a single transcript could result in one mRNA and the other without the 87 bases. In support of the latter hypothesis, we have shown that there is probably only one human dopamine D₂ receptor gene, DRD2, and that the 87-bp sequence is contained on a distinct exon of that gene. Furthermore, we have cloned a rat brain cDNA that contains the 87-bp sequence (unpublished results). This sequence is highly similar to that of the human cDNA and established that dopamine D₂ receptors containing the 29 residues are not unique to the human pituitary.

The human dopamine D₂ receptor expressed in L-hpitD₂Zem cells has essentially the same drug binding profile as do rat striatum and L-RGB2Zem-1 membranes. Therefore, since the 29 amino acids do not affect binding, they may be involved in other levels of receptor function. For example, the third cytoplasmic loop of the β_2 -adrenergic receptor has been shown to be required for appropriate G-protein coupling (21). Therefore, one possibility is that this sequence may influence whether dopamine D₂ receptor stimulation inhibits adenylyl cyclase, activates potassium channel conductance or inhibits calcium mobilization (22). Another possibility is that the 29 amino acids differentiate post-synaptic dopamine D₂ receptors from presynaptic autoreceptors (23). Since a computer search (VAX/Intelligenetics) failed to identify another sequence of significant homology, we consider this sequence to be unique to the D₂ receptor.

Based on the comparison of genomic and cDNA sequences, the human dopamine D₂ receptor gene is divided into at least seven exons. It is possible that one or more additional exons remain to be identified at the 5' end of the gene, as was shown to be the case with muscarinic receptor genes (24).

The interruption of coding sequence by introns distinguishes the human dopamine D₂ receptor gene from most other members of the G-protein coupled receptor gene family with the exception of the opsin genes (25). One significant observation is that the placement of two introns (Nos. 3 and 5) in this human gene corresponds almost precisely to intron positions conserved in bovine and Drosophila opsin genes (26, 27). The simplest interpretation of this finding is that their common ancestor, a gene roughly one billion years old (28), contained these introns. Our characterization of the gene structure also provides evidence that the exons encode recognizable elements of protein structure (29). That introns are found following transmembrane segments II, III, and IV (See Figs. 18 and 21) argues that the repeated structural motifs characteristic of these receptors may have evolved by internal duplication. Furthermore, the presence of several introns within the third cytoplasmic loop provides an explanation of the substantial variation in length observed across the family (30).

The possibility of alternatively spliced dopamine D₂ receptor mRNAs giving rise to structurally distinct forms is exciting. The expression of one form of the dopamine D₂ receptor mRNA or another represents a level of control which may have implications with respect to human disease.

Acknowledgements

We would like to thank Howard Goodman for discussion and review of the manuscript, Dee Yarozeski for manuscript preparation, and Vicky Robertson, Nancy Kurkinen, and June Shiigi for the illustrations. D.K.G. holds a fellowship from NIH. This work was supported by NIH Grant Nos. Dk37231 and MH45614 and a grant from Cambridge NeuroScience Research, Inc., Cambridge, Massachusetts, to O.C.

5

References

1. Lee, T., Seeman, P., Rajput, A., Farley, I.J. and
Hornykiewicz, O. (1978) *Nature* 273, 59-61.
2. Seeman, P., Ulpian, G., Bergeron, C., Riederer, P.,
Jellinger, K., Gabriel, E., Reynolds, G.P. and
Tourtellotte, W.W. (1984) *Science* 225, 728-730.
3. Kebabian, J.W. (1978) *Adv. Biochem.
Psychopharmacol.* 19, 131-154.
4. DeCamilli, P., Maconi, D. and Spada, A. (1979)
Nature (London) 278, 252-255.
5. Clark, D. and White, F.J. (1987) *Synapse* 1, 347-
388.
6. Weiner, R.I. and Ganong, W. (1978) *Physiol. Rev.*
58, 905-976.
7. Seeman, P. and Lee, T. (1975) *Science* 188, 1217-1219.
8. Creese, I., Burt, D.R., and Snyder, S.H. (1976)
Science 192, 481-483.
9. Bunzow, J.R., Van Tol, H.H.M., Grandy, D.K., Albert,
P.R., Salon, J., Christie, M., Machida, C.A.,
Neve, K., and Civelli, O. (1988) *Nature* 336,
783-787.
10. Gorman, C., Padmanabhan, R. and Howard, B.H. (1983)
Science 221, 551-553.
11. Albert, P.R., Neve, K., Bunzow, J.R. and Civelli, O.
(1989) submitted.
12. Grigoriadis, D. and Seeman, P. (1986) *Naunyn-
Schmiederberg Arch. Pharmacol.* 321, 21-25.
13. Baudry, M., Martres, M.P. and Schwartz, J.C. (1979)
Naunyn-Schiederberg Arch. Pharmacol. 308, 231-
237.

14. Fischer, J.B. and Schonbrunn, A. (1988) J. Biol. Chem. 263 2808-2816
15. Grandy, D.K. and Dodgson, J.B. (1987) Nuc. Acids Res. 15, 1063-1080.
- 5 16. Church, G.M. and Gilbert, W. (1984) Proc. Natl. Acad. Sci. USA 81, 1991-1995.
17. Marchionni, M. and Gilbert, M. (1986) Cell 46, 133-141.
18. O'Dowd, B.F., Hnatowich, M., M.C.G., Lefkowitz, R.J. and Bouvier, M. (1989) J. Biol. Chem. 264, 7564-7569.
- 10 19. Grandy, D.K., Litt, M., Allen, L., Bunzow, J.R., Marchionni, M.A., Makam, H., Frothingham, L., Magenis, R.E. and Civelli, O. (1989) Am. J. Hum. Genet. accepted.
- 15 20. Mount, S.M. (1982) Nuc. Acids Res. 10, 461-472.
21. Strader, C.D., Dixon, R.A.F., Cheung, A.H., Candelore, M.R., Blake, A.D. and Sigal, I.S. (1987) J. Biol. Chem. 262, 16439-16443.
- 20 22. Vallar, L. and Meldolesi, J. (1989) TIPS 10, 74-77.
23. Tepper, J.M., Gariano, R.F. and Groves, P.M. (1987) in Neurophysiology of Dopaminergic Systems: Current Status and Clinical Perspectives, eds. Chiodo, L.A. and Freeman, A.S. (Lakeshore Publishing Co., Michigan), pp. 93-127.
- 25 24. Peralta, E.G., Winslow, J.W., Peterson, F.L., Smith, D.H., Ashkenazi, Avi, Ramachandran, J., Schimerlik, M.I. and Capon, D.J. (1987) Science 236, 600-605.
- 30 25. O'Dowd, B.F., Lefkowitz, R.J. and Caron, M.G. (1989) Ann. Rev. Neurosci. 12, 67-83.
26. Nathans, J. and Hogness, D.S. (1983) Cell 32, 807-814.
- 35 27. O'Tousa, J.E., Baehr, W., Martin, R.L., Hirsch, J., Pak, W.L. and Applebury, M.L. (1985) Cell 40, 839-850.

28. Doolittle, R.F. (1987) Of URFS and ORFS (University Science Books, Mill Valley), p. 40.
29. Gilbert, W. (1978) Nature 271, 501.
30. Dohlman, H.G., Caron, M.G. and Lefkowitz, R.J.
(1987) Biochem. 26, 2657-2664.

5

WHAT IS CLAIMED IS:

1. An isolated DNA sequence of Claim 53 encoding a polypeptide having the biological activity of a mammalian D₂ dopamine receptor, or a nucleic acid sequence useful as a probe for and which hybridizes with such a sequence.
2. A DNA sequence of Claim 53 having the sequence recited in Figure 1, a complementary strand thereof, a sequence differing therefrom by codon degeneracy, a sequence which hybridizes therewith and encodes a mammalian D₂ dopamine receptor, or a nucleic acid sequence useful as a probe for and which hybridizes with such a sequence.
3. An isolated DNA sequence encoding a polypeptide having the biological activity of a dopamine receptor.
4. A nucleic acid sequence of Claim 53 which is a mammalian D₂ dopamine receptor gene or a fragment thereof coding for a polypeptide having the biological activity of such a receptor.
5. A nucleic acid sequence of Claim 53, wherein said mammal is human.
6. A DNA sequence of Figure 7, a complementary strand thereof, a sequence differing therefrom by codon degeneracy, a sequence which hybridizes therewith and encodes a human D₂ dopamine receptor, a nucleic acid sequence useful as a probe for and which hybridizes with such a sequence, or a sequence which comprises one of said sequences.

7. An oligonucleotide useful as a probe which hybridizes with a nucleic acid sequence of Claim 53.

8. An oligonucleotide useful as a probe which hybridizes with a nucleic acid sequence of Claim 1.

9. An oligonucleotide useful as a probe which hybridizes with a nucleic acid sequence of Claim 5.

10. An oligonucleotide probe of Claim 7 labelled with a detectable moiety.

11. A oligonucleotide probe of Claim 7 encoding a mammalian D₂ dopamine receptor fragment from one of its seven transmembrane domains or from its third cytoplasmic loop.

12. A vector comprising DNA of Claim 53.

13. A vector comprising DNA of Claim 1.

14. A vector of Claim 12, wherein the DNA is under the control of a viral promoter.

15. A vector of Claim 12, which is vaccinia virus or a retrovirus.

16. A cell transformed with a vector of Claim 12.

17. A cell transformed with a vector of Claim 13.

18. A cell transformed with a vector of Claim 12, wherein said mammal is human.

19. A cell of Claim 16 from a multicellular organism.

20. A cell of Claim 19 which is a mammalian cell.

21. A method for preparing a polypeptide having the biological activity of a dopamine receptor comprising culturing a cell of Claim 16.

22. A method for preparing a polypeptide having the biological activity of a dopamine receptor comprising culturing a cell of Claim 18.

23. A method of probing a nucleic acid sample for the presence of a sequence which hybridizes with a sequence encoding a neurotransmitter receptor, comprising hybridizing said sample with the 1.3 kb HindIII fragment of the hamster β_2 AR gene or with a probe which hybridizes with said fragment.

24. A substantially pure polypeptide of Claim 29 which is a mammalian D_2 dopamine receptor.

25. A receptor of Claim 24 encoded by a DNA sequence of Figure 1 or a sequence which hybridizes therewith.

26. A receptor of Claim 24 wherein said mammal is human.

27. A substantially pure mammalian D_2 dopamine receptor, wherein at least one amino acid is substituted for, inserted into and/or deleted from a dopamine receptor of Claim 24.

28. A dopamine receptor of Claim 27 which is a mutant having the biological activity of a dopamine receptor.

29. A substantially pure polypeptide having the biological activity of a mammalian D₂ dopamine receptor.

30. A polypeptide of Claim 29 which is a fragment of a human D₂ dopamine receptor.

31. A fragment of a mammalian D₂ dopamine receptor having an amino acid sequence which defines an antigenic determinant of said receptor.

32. A fragment of Claim 31 selected from the third cytoplasmic loop of the receptor.

33. An antibody capable of binding a mammalian D₂ dopamine receptor.

34. A monoclonal antibody of Claim 33.

35. A fragment of an antibody of Claim 33, which fragment is capable of binding a dopamine receptor.

36. An antibody fragment of Claim 35 which is the Fab fragment.

37. An antibody of Claim 33, wherein said mammal is human.

38. An antibody of Claim 33 which is labelled.

39. A composition comprising a polypeptide having the biological activity of a mammalian D₂ dopamine receptor or a biologically active fragment or variant thereof, an antibody or antibody fragment thereof, or a nucleic acid sequence of Claim 1, and a pharmaceutically acceptable carrier.

40. A composition of Claim 39 wherein said mammal is human.

41. A method of increasing the mammalian D₂ dopamine receptor activity at a locus, comprising placing at the locus, or causing to be expressed at the locus, an isolated DNA sequence of Claim 1 encoding a polypeptide having the biological activity of a mammalian D₂ dopamine receptor, or placing at the locus such a polypeptide in substantially pure form.

42. In a method of determining the effect of a condition or substance on a mammalian D₂ dopamine receptor comprising exposing such a dopamine receptor to said condition or substance, the improvement wherein a polypeptide of Claim 29 is used as said dopamine receptor.

43. A method of Claim 42 comprising screening a chemical compound for affinity to a mammalian D₂ dopamine receptor or for its ability to modify or initiate a response triggered by ligand binding to a mammalian D₂ dopamine receptor.

44. An antibody of Claim 33 which binds a partial sequence of Figure 1, which comprises amino acids numbers 2-13, 182-192, 264-277, 287-298 or 404-414.

45. A transgenic nonhuman mammal in which a nucleic acid sequence of Claim 53 is integrated into its chromosome at a locus whereby the sequence is active as a mammalian D₂ dopamine receptor gene.

46. A method of determining a dopamine receptor-related property of a sample, comprising exposing at least a component of said sample to a nucleic acid or

amino acid sequence uniquely associated with a dopamine receptor.

47. A method of claim 46, wherein said sample is from a human and a disease state associated with dopamine receptors is diagnosed or characterized.

48. A method of claim 47, wherein said disease state is schizophrenia.

49. A polypeptide having the biological activity of a mammalian D₂ dopamine receptor, which is prepared by expression in a cell of Claim 16.

50. A polypeptide having the biological activity of a mammalian D₂ dopamine receptor, which is prepared by expression in a cell of Claim 17.

51. A polypeptide having the biological activity of a mammalian D₂ dopamine receptor, which is prepared by expression in a cell transformed with a vector comprising DNA of Claim 5.

52. A method of modulating mammalian D₂ dopamine receptor activity, comprising administering a composition of Claim 54.

53. A mammalian D₂ dopamine receptor gene in substantially pure form or a biologically active fragment thereof.

54. A composition comprising a polypeptide having the biological activity of a mammalian D₂ dopamine receptor or a biologically active fragment or variant thereof, an antibody or antibody fragment thereof, or a nucleic acid sequence of Claim 53, and a pharmaceutically acceptable carrier.

1 / 47

FIG. 1-1

GGCTGCCGG

-120

AGGGGGGGCGTGCGTGATGCGGGGAGCTGGAAGCCTCGAGCAGCCGGCCCTTCT

CTGGCCCCGGGCCATATGGCTTGAAGAGCCGTGCCACCCAGTGGCCCCACTGCCCA
-1

1	Met	Asp	Pro	Leu	Asn	Leu	Ser	Trp	Tyr	Asp	Asp	Leu	Glu	Arg
	ATG	GAT	CCA	CTG	AAC	CTG	TCC	TGG	TAC	GAT	GAC	CTG	GAG	AGG

1	Gln	Asn	Trp	Ser	Arg	Pro	Phe	Asn	Gly	Ser	Glu	Gly	Lys	Ala	Asp
	CAG	AAC	TGG	AGC	CGG	CCC	TTC	AAT	GGG	TCA	GAA	GGG	AAG	GCA	GAC

	Arg	Pro	His	Tyr	Asn	Tyr	Tyr	Ala	Met	Leu	Leu	Thr	Leu	Leu	Ile
	AGG	CCC	CAC	TAC	AAC	TAC	TAT	GCC	ATG	CTG	CTC	ACC	CTC	CTC	ATC

	Phe	Ile	Ile	Val	Phe	Gly	Asn	Val	Leu	Val	Cys	Met	Ala	Val	Ser
	TTT	ATC	ATC	GTC	TTT	GGC	AAT	GTG	CTG	GTG	TGC	ATG	GCT	GTA	TCC

2 / 47

Arg	Glu	Lys	Ala	Leu	Gln	Thr	Thr	Thr	Asn	Tyr	Leu	Ile	Val	Ser	70
GCA	GAG	AAG	GCT	TTG	CAG	ACC	ACC	ACC	AAC	TAC	TTG	ATA	GTC	AGC	
Leu	Ala	Val	Ala	Asp	Leu	Leu	Val	Ala	Thr	Leu	Val	Met	Pro	Trp	90
CTT	GCT	GTG	GCT	GAT	CTT	CTG	GTG	GCC	ACA	CTG	GTA	ATG	CCG	TGG	
Val	Val	Tyr	Leu	Glu	Val	Val	Gly	Glu	Trp	Lys	Phe	Ser	Arg	Ile	100
GTT	GTC	TAC	CTG	GAG	GTG	GTG	GGT	GAG	TGG	AAA	TTC	AGC	AGG	ATT	
His	Cys	Asp	Ile	Phe	Val	Thr	Leu	Asp	Val	Met	Met	Cys	Thr	Ala	120
CAC	TGT	GAC	ATC	TTT	GTC	ACT	CTG	GAT	GTC	ATG	ATG	TGC	ACA	GCA	
Ser	Ile	Leu	Asn	Leu	Cys	Ala	Ile	Ser	Ile	Asp	Arg	Tyr	Thr	Ala	130
AGC	ATC	CTG	AAC	CTG	TGT	GCC	ATC	AGC	ATT	GAC	AGG	TAC	ACA	GCT	
Val	Ala	Met	Pro	Met	Leu	Tyr	Asn	Thr	Arg	Tyr	Ser	Ser	Lys	Arg	150
GTG	GCA	ATG	CCC	ATG	CTG	TAT	AAC	ACA	CGC	TAC	AGC	TCC	AAG	CGC	
															450

FIG.1-2

3 / 47

Arg	Val	Thr	Val	Met	Ile	Ala	Ile	Val	Trp	Val	Leu	Ser	Phe	Thr
CGA	GTT	ACT	GTC	ATG	ATT	GCC	ATT	GTC	TGG	GTC	CTG	TCC	TTC	ACC
									160					
Ile	Ser	Cys	Pro	Leu	Leu	Phe	Gly	Leu	Asn	Asn	Thr	Asp	Gln	Asn
ATC	TCC	TGC	CCA	CTG	CTC	TTC	GGA	CTC	AAC	AAT	ACA	GAC	CAG	AAT
														540
Glu	Cys	Ile	Ile	Ala	Asn	Pro	Ala	Phe	Val	Val	Tyr	Ser	Ser	Ile
GAG	TGT	ATC	ATT	GCC	AAC	CCT	GCC	TTT	GTG	GTC	TAC	TCC	TCC	ATT
									190					
Val	Ser	Phe	Tyr	Val	Pro	Phe	Ile	Val	Thr	Leu	Leu	Val	Tyr	Ile
GTC	TCA	TTC	TAC	GTG	CCC	TTC	ATC	GTC	ACT	CTG	CTG	GTC	TAT	ATC
														630
Lys	Ile	Tyr	Ile	Val	Leu	Arg	Lys	Arg	Arg	Lys	Arg	Val	Asn	Thr
AAA	ATC	TAC	ATC	GTC	CTC	CGG	AAG	CGC	CGG	AAG	CGG	GTC	AAC	ACC
									220					
Lys	Arg	<u>Ser</u>	<u>Ser</u>	Arg	Ala	Phe	Arg	Ala	Asn	Leu	Lys	Thr	Pro	Leu
AAG	CGC	AGC	AGT	CGA	GCT	TTC	AGA	GCC	AAC	CTG	AAG	ACA	CCA	CTC
														720

FIG. 1-3

SUBSTITUTE SHEET

4 / 47

```

Lys Asp Ala Ala Arg Arg Ala Gln Glu Leu Glu Met Glu Met Leu
AAG GAT GCT GCC GCC CGC CGA GCT CAG GAG CTG GAA ATG GAG ATG CTG
250

Ser Ser Thr Ser Pro Pro Glu Arg Thr Arg Tyr Ser Pro Ile Pro
TCA AGC ACC AGC CCC CCA GAG AGG ACC CGG TAT AGC CCC ATC CCT
260 270

Pro Ser His His Gln Leu Thr Leu Pro Asp Pro Ser His His Gly
CCC AGT CAC CAC CAG CTC ACT CTC CCT CCT GAT CCA CCA TCC CAC CAC GGC
280

Leu His Ser Asn Pro Asp Ser Pro Ala Lys Pro Gly Lys Asn Gly
CTA CAT AGC AAC CCT GAC GAC AGT CCT GGC GCC AAA CCA CCA GAG AAG AAT GGC
290 300

His Ala Lys Ile Val Asn Pro Arg Ile Ala Lys Phe Phe Glu Ile
CAC GCC AAG ATT GTC AAT AAT CCC AGG ATT GCC AAG TTC TTC TTT GAG ATC
310

Gln Thr Met Pro Asn Gly Lys Thr Arg Thr Ser Leu Lys Thr Met
CAG ACC ATG CCC AAT GGC AAA ACC CGG ACC TCC CTT AAG ACG ATC
320 330

```

FIG. 1-4

5 / 47

```

Ser Arg Arg Lys Lys Leu Ser Gln Gln Lys Lys Lys Ala Thr Gln
AGC CGC AGA AAG CTC TCC CAG CAG AAG AAG AAA GCC ACT CAG
340

Met Leu Ala Ile Val Leu Gly Val Phe Ile Ile Cys Trp Leu Pro
ATG CTT GCC ATT GTT CTC CTC GGT GTG TTC ATC ATC TGC TGG CTG CCC
350
1080
Phe Phe Ile Thr His Ile Leu Asn Ile His Cys Asp Cys Asn Ile
TTC TTC ATC ACG CAC CAC ATC CTG AAT ATA CAC TGT GAT TGC AAC ATC
370

Pro Pro Val Leu Tyr Ser Ala Phe Thr Trp Trp Leu Gly Tyr Val Asn
CCA CCA GTC CTC TAC AGC AGC GCC TTC ACA TGG CTG GGC TAT GTC AAC
380
1170
Ser Ala Val Asn Pro Ile Ile Ile Tyr Thr Thr Phe Asn Ile Glu Phe
AGT GCC GTC AAC CCC ATC ATC ATC ATC ACC ACC ACC TTC AAC ATC GAG TTC
400

Arg Lys Ala Phe Met Lys Ile Leu His Cys
CGC AAG GCC TTC ATG AAG AAG ATC TTG CAC TGC TGAGTCTGCCCTTGCCCTG
410
1264

```

FIG.1-5

SUBSTITUTE SHEET

6 / 47

CACAGCAGCTGCTTCCACCTCCCTGCCTATGCAGGCCAGACCTCATCCCTGCAAGCTG
TGGGCAGAAAGCCCAGATGAACCTTGGCCTTCTCTCGACCCCTGCAGGCCCTGCAGTGTTA
GCTTGGCTCGATGCCCTCTCTGCCCACACACCCCTCATCTGCCAGGGTAGGGCCAGGG 1383
AGACTGGTATCTTACCAGCTCTGGGGTTGGACCCATGGCTCAGGGCAGCTCACAGAGTGC
CCCTCTCATATCCAGACCCCTGTCTCCTTGGCACCAAGATGCAGCGGCCTTCCTTGACC 1502
TTCCCTCTTGGGCACAGAAACTAGCTCAGTGTCTCGAGCACACCCCTGATCGCTGGCTTGCCC
TGGCCCTTGCTTGCCCTGTGCCGGATCAGGTGGTGGGAGGAGCGACACGTTCTTACTTT 1621
ATAGGAACCACATAGGAAAGCAGGGAAACACGCCAAGTCTCCAGGCACATCAGTGTCAAG 1740
AGACACACATAAACACACAGGTAGCTCCATGGACCCCCAGAGAAACTGAGGCTGAAAAATC
TGTTTCCACTCCAACTCTAGTGTGAGTCCCTACTTTTCATAGCCATGGGTATTACTATG 1859

FIG. 1-6

7 / 47

TCCTACCTTGTTATAGTATCCCATGGGTTTCTGTACCATTTGGGGGAAACAACACTCTA
ATCCTCAAGGGCCCAAGAGAATCTGTAAGGAGAAAAATAGGCTGATCTCCCTCTACTCT
1978
CCAATCCACTCCACCACCTTCTTGATATACCTTGGATGTATCCATTCCCTCACAGCAAAATG
CTGGCCAGTCAGGCCCTTGACCAGTGTGGAGTTGAAGCTGGATGTGGTAACCTTGGGGCT
2097
CTTTGGGCTGGGGGTTGTAAACATCGTCTCTCTTCCATATCTCTTCCCTTCCAGTG
CCTCTGCCCTTAGAAGAGGCTGTGGATGGGGTGGTGGACTGCTGATACCATTTGGGCCCTGG
2216
CCCTGAATGAGGAGGGGAAGCTGCAGTTTGGAGGGTTCTGGGATCCAACCTCTGTAAACAT
CACTATACCTGTACCAAACTAATAAAACCTTGACAAGAGTCAAAAAA
2317

FIG.1-7

8 / 47

FIG. 2

I

MD...[FL...NISWYDDDLERQNSRPFNGSEKADRPYNYAMITLL].IFIIVFGNVLVCMAVSREKALQITINY
MGP...[P...CNDSDFLLTNGSHV...PDHDVTEERDEAWVCGMAIMSVIVLAVFGNVLVITAIKFERLQITINY
MGSQ...[POA.CNASWNGTEAPG..GGARATPYSLOVT...LTLVCLAGLL.MLLTVFGNVLVIIVFTSRATKAPQNL
MDVLS.[EGQ.QNNTTSPPAFFE..TCGNTTGISDVTVSQV.ITSLITGIL].IFCAVTCGACVVAJALERSLONVANY
MNTSAPPAVSPNITVLAP.....GKGPWQVA.....FIGITGIL.SLATVTCGNLVLISFKVNTLKITVNNY
MGACV.VMTDINIS.....SGLDSNATGITAFSMPGWQLALWTAAYLAL.VLVAVMGNATVILWILAHQRMRTIVINY

III

II

LIVSLAMADIIIVAILVMPWVYLEVGVGEMKFSRIHCDIFVTLDVMMCTASILNLCASISQRYTAVAMPIMTRYSSKRR
FIDSLACADLVGIAVVPFGASHIIMKVMNFGNFWCEFTSILVLCVITASIELCVIAVQRYIAITSEFKYOSLLTKNKA
FIVSLASADILIVATLVIPFSLANEMQVMEGKTWCEIYALDVLCTSSIVHLCAISQRYMSITQAIENLKRTPRR
LIGSLAVTDIMVSVLVIPMAALYQVLMNMLGQVTCDFIALDVLCTSSILHLCAIAQRYWATIDFIDVYNKRTPPR
FLLSLACADILIGFMSNLYTTLMLQVMAIGTACDLWALDYVASNASUMNLLIISQRYFSVTRPLSYRAKTRPRA
FLVNLALADICMAAFNAAFNFVYASHNIMVFGRAFQYQNLFPITAMFVSIYSMTAJAAQRYMAIVRPFQRLSAPGTR.

IV

V

VTW[TAIVWVISETISC.[PLLLGLNNTD.....QNECIANPAFVVYSSIMSFYVPEFIVTLVYIKIVLVRKRKRVTNR--(111)
RMV.[ILMVWLVSGLTISFIPIQMHWYRATH..QKAI DCYHRETCDDFTNQAYATMSSISVSFVPIVMMVVFVYSRVFQVAKRQLKI--(32)-
KAI.[ITVWVISAVISFPLISIEKKGGG.....GGPQPAEPRCEINDQKWVHSSCIGSFAPOHIMILVYRIYQIAKERTRPV--(137)
ALT.SLT.MLIGFLISIPMLGWRTPEDR.....SDPDACTISKDMGYTUYSTFGAFYIPILMLVLYGRIFRAARFIPIT--(110)
ALM.[GLAWLVSEVLWA.[PAILFWQYLVE.....RTVLACQCYIQFLSQPIITFGTAMAAFLPVTIMCTIIMRIYRETERAREL--(137)
.AV.[IAGIWLVALALAF.[EQCFYSTITDEGATKCVVANPEDSGGKMILLYHLIVIALIYE.LPVMVFAVSIGLTLWRSVPG--(12)-

VI

VII

-KEKKATQMTAIVIGMFILCWLPPFFITHIINIHCOCN.....IPPVLYSAFTWLGYNVA..VNPIIYITFENIEFRNAFMKIIHC
-KEHKALKTIGIIMGIFILCWLPPFFIVNIVHVIQDNL.....IPKEVYILLNWLGYVNSA..PNPIIYCRSP.DFRIAFQELI..CL--(36)
-REKRTFVIAVILGMFVVCWEPFFETYLIAVGCSS.....VPRTEFKFFWEGYCNSS..INPVIYTLFNHDFRFAFMKII.CRC--(6)
-RERTVKTLOIIMGIFILCWLPPFFIVALVLPFCSSC..HMPITLGAIIINWLGYNSL..INPVIYAYENKDFQNAFMKIIKCNFCRQ
-KEKKAARTLSAILLAFIVIMV[YNIMVLVSTFCCKDC.....VPEITIMELGYWIGYVNST..INPMCYALQNKAFRDTFRLIIFCR--(24)
-AKKRFVKMTMIVVVTFAICWLPHYLYHIGTFQEDIYCHKFIQVYLALPWIA..MSSTMYNPIIYCCINHRFSQERLAPRC--(62)

9 / 47




FIG. 3

10 / 47




FIG.4A-1





FIG.4A-2

11 / 47

FIG. 4B



SUBSTITUTE SHEET

12 / 47

DRUG

Ki (nM)

RGB-2

Transformed Ltk-Cells

Rat Striatum


(+)-Butaclamol	0.83	1.0
(-)-Butaclamol	>1,000	>1,000
Haloperidol	3.0	5.3
Dopamine + GTP	17,000	6,300
Sulpiride	80	67 (87%)
high affinity	---	>10,000 (13%)
low affinity	---	35 (16%)
SCH 23390	1,000	780 (84%)
high affinity	---	27 (25%)
low affinity	---	>1,000 (75%)
Ketanserin	---	
high affinity	---	
low affinity	>1,000	

SUBSTITUTE SHEET


SUBSTITUTE SHEET

FIG. 4C

13 / 47



RGB-2




SUBSTITUTE SHEET

14 / 47

FIG. 6-1

Length of DNA: 2477 nts

**SUBSTITUTE SHEET**

15 / 47





FIG.6-2

SUBSTITUTE SHEET

16 / 47



SUBSTITUTE SHEET

FIG.6-3

17 / 47




FIG. 6-4

SUBSTITUTE SHEET

18 / 47




FIG. 6-5

FIG. 7-1

3' GTC GGG TGG GCT CTC CTG GGC CAT GTC GGG GTA GGG TGG GTC GGT GGT GGT CGA 5'
 5' CAG CCC ACC CGA GAG GAC CCG GTA CAG CCC CAT CCC ACC CAG CCA CCA CCA GCT 3'
 Gln Pro Thr Arg Glu Asp Pro Val Gln Pro His Pro Thr Gln Pro Pro Pro Ala
 Ser Pro Pro Glu Arg Thr Arg Tyr Ser Pro Ile Pro Pro Ser His His Gln Leu
 Ala His Pro Arg Gly Pro Gly Thr Ala Pro Ser His Pro Ala Thr Thr Ser

81
 CTG AGA GGG GCT GGG CAG GGT GGT ACC AGA GGT GTC GTG AGG GCT GTC GGG GCG
 GAC TCT CCC CGA CCC GTC CCA CCA TGG TCT CCA CAG CAC TCC CGA CAG CCC CGC
 Asp Ser Pro Arg Pro Val Pro Pro Trp Ser Pro Gln His Ser Arg Gln Pro Arg
 Thr Leu Pro Asp Pro Ser His His Gly Leu His Ser Thr Pro Asp Ser Pro Ala
 Leu Ser Pro Thr Arg Pro Thr MET Val Ser Thr Ala Leu Pro Thr Ala Pro Pro

135
 GTT TGG TCT CTT CTT ACC CGT ACG GTT TTC CTG GTG GGG TTC CTA ACG GTT CTA
 CAA ACC AGA GAA GAA TGG GCA TGC CAA AAG GAC CAC CCC AAG GAT TGC CAA GAT
 Gln Thr Arg Glu Glu Trp Ala Cys Gln Lys Asp His Pro Lys Asp Cys Gln Asp
 Lys Pro Glu Lys Asn Gly His Ala Lys Arg Thr Thr Pro Arg Ile Ala Lys Ile
 Asn Gln Arg Arg MET Gly MET Pro Lys Gly Pro Pro Gln Gly Leu Pro Arg Ser

189
 GAA ACT CTA GGT CTG GTA CCG GTT ACC GTT TTG GGC CTG GAG GGA GTT CTG GTA
 CTT TGA GAT CCA GAC CAT GCC CAA TGG CAA AAC CCG GAC CTC CCT CAA GAC CAT
 Leu . Asp Pro Asp His Ala Gln Trp Gln Asn Pro Asp Leu Pro Gln Asp His
 Phe Glu Ile Gln Thr MET Pro Asn Gly Lys Thr Arg Thr Ser Leu Lys Thr MET

SUBSTITUTE SHEET

FIG. 7-2

CTC GGC ATC CTT C6A G55 GGT CGT CTT CCT CTT TCG GTG AGT CTA CGA GCG 270
 GAG CCG TAG GAA G3T C44 CCA GCA GAA GAA GAG CAC TCA GAT GCT CGC
 Glu Pro . Glu C TT Pro Ala Glu Gly Glu Ser His Ser Asp Ala Arg
 Ser Arg Arg Lys Leu Phe Gln Gln Lys Glu Lys Ala Thr Gln MET Leu Ala
 Ala Val Gly Ser Ser Arg Arg Arg Lys Pro Leu Arg Cys Ser Pro
 243
 297
 GTA 5CA AGA GCC GCA CAA GTA GTA GAC CGA CGG GAA GAA GTA GTG TGT GTA 324
 CAT 4GT TCT CCG CGT GTT CAT CAT CTG CTG GCT GCC CTT CTT CAT CAC ACA CAT
 His T Ser Arg Arg Val His His Leu Leu Ala Ala Leu Leu His His Thr His
 Ile Val Leu Gly Val Phe Ile Ile Cys Trp Leu Pro Phe Phe Ile Thr His Ile
 le Phe Ser Ala Cys Ser Ser Ser Ala Gly Cys Pro Ser Ser His Thr Ser
 351
 378
 GGA CTT GTA TGT GAC ACT GAC GAC GGT GTA GGG CGG ACA GGA CAT GTC GCG GAA GTG
 CCT GAA CAT ACA CTG TGA CTG CAA CAT CCC GCC TGT CCT GTA CAG CGC CTT CAC
 Pro Glu His Thr Leu . Leu Gln His Pro Ala Cys Pro Val Gln Arg Leu His
 Leu Asn Ile His Cys Asp Cys Asn Ile Pro Pro Val Leu Tyr Ser Ala Phe Thr
 . Thr Tyr Thr Val Thr Ala Thr Ser Arg Leu Ser Cys Thr Ala Pro Ser Arg
 405
 432
 CAC CGA CCC GAT ACA GTT GTC GCG GCA CTT GGG GTA GTA GAT GTG GTG GAA GTT
 GTG GCT GCG CTA TGT CAA CAG CGC CGT GAA CCC CAT CAT CTA CAC CAC CTT CAA
 Val Ala Ala Gly Leu Cys Gln Gln Arg Arg Glu Pro His His Leu His His Leu Gln
 Trp Leu Gly Tyr Val Asn Ser Ala Val Asn Pro Ile Ile Tyr Thr Thr Phe Asn
 Gly Trp Ala MET Ser Thr Ala Pro . Thr Pro Ser Ser Thr Pro Pro Ser Thr

SUBSTITUTE SHEET

459
 GTA ACT CAA GGC GTT CCG GAA GGA CTT CTA GGA GGT GAC GAC TGA GAC GAC GGA 486
 CAT TGA GTT CCG CAA GGC CTT CCT GAA GAT CCT CCA CTG ACT CTG CTG CCT
 His . Val Pro Gln Gly Leu Pro Glu Asp Pro Leu Leu Thr Leu Leu Pro
 Ile Glu Phe Arg Lys Ala Phe Leu Lys Ile Leu His Cys STOPLeu Cys Cys Leu
 Leu Ser Ser Ala Arg Pro Ser . Arg Ser Ser Thr Ala Asp Ser Ala Ala Cys
 513
 CGG CGT GTC GTC GGA CGA AGG GTG GAG GGA CGG GTC ACG GCC GGT CGG AGT GGG 540
 GCC GCA CAG CAG CCT GCT TCC CAC CTC CCT GCC CAG TGC CGG CCA GCC TCA CCC
 Ala Ala Gln Gln Pro Ala Ser His Leu Pro Ala Gln Cys Arg Pro Ala Ser Pro
 Pro His Ser Ser Leu Leu Pro Thr Ser Leu Pro Ser Ala Gly Gln Pro His Pro
 Arg Thr Ala Ala Cys Phe Pro Pro Cys Pro Val Pro Ala Ser Leu Thr Leu
 567
 AAC GCT TGG CAC TCG TCC TTC CGG ACC CAC CTA GCC GGA GGA GAA GAT CGG GGC 594
 TTG CGA ACC GTG AGC AGG AAG GCC TGG GTG GAT CGG CCT CCT CTT CTA GCC CCG
 Leu Arg Thr Val Ser Arg Lys Ala Trp Val Asp Arg Pro Pro Leu Leu Ala Pro
 Cys Glu Pro . Ala Gly Arg Pro Gly Trp Ile Gly Leu Leu Phe . Pro Arg
 Ala Asn Arg Glu Gln Glu Gly Leu Gly Ser Ala Ser Ser Ser Pro Gly

21/47

FIG. 7-3

SUBSTITUTE SHEET

22 / 47





FIG. 8



23 / 47

FIG. 9A







FIG. 9B



24 / 47



25 / 47

CONTROL				+P.T.			
BASAL	FSK	FSK+DA		BASAL	FSK	FSK+DA	
2.1	22.6	14.3		1.7	26.7	23.1	
0.2	2.2	2.1		0.4	2.0	0.5	
—	—	41%		—	—	14%	

\bar{X}
S.E.
INH

FIG. 12A

CONTROL				+P.T.			
BASAL	VIP	DA	VIP+DA	BASAL	VIP	DA	VIP+DA
0.60	2.41	0.32	0.84	0.61	2.68	0.55	2.56
0.02	0.31	0.02	0.13	0.09	0.08	0.03	0.25
—	—	53%	71%	—	—	10%	3%

\bar{X}
S.E.
INH

FIG. 12B

CONTROL				+P.T.			
BASAL	VIP	DA	VIP+DA	BASAL	VIP	DA	VIP+DA
0.78	5.1	0.25	0.76	0.64	5.29	0.66	4.76
0.04	0.4	0.03	0.01	0.01	0.44	0.03	0.16
—	—	68%	88%	—	—	0%	12%

\bar{X}
S.E.
INH

FIG. 12C

26 / 47




FIG. 13 A

27 / 47

FIG. 13B-1





FIG. 13B-2



28 / 47

FIG. 13C





FIG. 14C

D

D(-S)

SUBSTITUTE SHEET

29 / 47

FIG.14A





FIG.14B



30 / 47

FIG. 15A






FIG. 15B



SUBSTITUTE SHEET



32 / 47

FIG.17A





FIG.17B



SUBSTITUTE SHEET

33 / 47

```

-33 AGAGCCTGGCCACCCAGTGGCTCCACCGCCCTG

      *           10           *           20
METAspProLeuAsnLeuSerTrpTyrAspAspLeuGluArgGlnAsnTrpSerArg
ATGGATCCACTGAATCTGTCCTGGTATGATGATCTGGAGAGGCAGAACTGGAGCCCG
|||||
ATGGATCCACTGAACCTGTCCTGGTACGATGACGATCTGGAGAGGCAGAACTGGAGCCCG
      *           30           *           40
ProPheAsnGlySerAspGlyLysAlaAspArgProHisTyrAsnTyrTyrAlaThrLeu
CCCTTCAACGGGTCAGACGGGAAGCGGACAGACCCCACTACTATGCCACACTG
|||||
CCCTTCAATGGGTCAGAAAGGGAAGGCAGACAGGCCCACTACTATGCCATGCTG
      Glu
MET
120

```

FIG. 18-I

SUBSTITUTE SHEET

34 / 47

240

[illegible]

FIG. 18-2

SUBSTITUTE SHEET

36 / 47

IV 170 180
ValLeuSerPheThrIleSerCysProLeuLeuPheGlyLeuAsnAsnAlaAspGlnAsn
 GTCCGTGTCCTTCACCATCTCCTGCCCACTCCTCTTCGGACTCAATAACGCAGACCAGAAC
 !!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!
 GTCCGTGTCCTTCACCATCTCCTGCCCACTGCTCTTCGGACTCAACAATACAGACCAGAAT
 Thr

190 V 200
GluCysIleIleAlaAsnProAlaPheValValTyrSerSerIleValSerPheTyrVal
 GAGTGTCATCATGCGCAACCCGGCCTTCGTGGTCTACTCCTCCATCGTCTCCTTCTACGTG
 !!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!! 600
 GAGTGTCATCATGCGCAACCCGGCCTTGTGGTCTACTCCTCCATTGTCCTCATTTCTACGTG

210 220
ProPheIleValThrLeuLeuValTyrIleLysIleTyrIleValLeuArgArgArgArg
 CCCTTCATTGTCACCCCTGCTGGTCTACATCAAGATCTACATTGTCTCTCCGCAGACGCCGC
 !!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!
 CCCTTCATCGTCACTGCTGCTATATCAAAATCTACATCGTCTCTCCGGAAGCGCCGG
 Lys

FIG. 18-4

SUBSTITUTE SHEET

37 / 47

LysArgValAsnThrLysArgSerArgAlaPheArgAlaHisLeuArgAlaProLeu 230 240
AAGCGAGTCAACACCAACGACGAGCCGAGCTTTCAGGGCCACCTGAGGGCTCCACTA
IIIIII IIIIIIIII IIIIIIIII III IIIIIII III IIIII 720
AAGCGGTCAACACCAAGCGCAGCAGTCGAGCTTTCAGAGCCAACTGAAGACACCACTC
Asn LysThr
* 250 *
LysGlyAsnCysThrHisProGluAspMETLysLeuCysThrValIleMETLysSerAsn
AAGGGCAACTGTACTCACCCTCGAGGACATGAAACTCTGCACCGTTATCATGAAGTCTAAT
III
AAG.....
GlySerPheProValAsnArgArgArgValGluAlaAlaArgArgAlaGlnGluLeuGlu 270 280
GGGAGTTTCCAGTGAACAGCGGAGAGTGGAGGCTGCCCGCGAGCCAGGAGCTGGAG
II IIIIIIIII IIIIIIIII IIIIIIIIIII
.....GATGCTGCCCGCGAGCTCAGGAGCTGGAA 840
Asp

FIG. 18-5

38 / 47

290 300
 METGluMETLeuSerSerThrSerProGluArgThrArgTyrSerProIleProPro
 ATGGAGATGCTCTCCAGCACCGCCACCCGAGAGAGACCCGGTACAGCCCATCCACCCC
 IIIIIIIII IIIIIIIIIII IIIIIIIIIII IIIIIIIIIII IIIIIIIIIII IIIIIIIIIII
 ATGGAGATGCTGTCAAGCACCGCCCGCCAGAGAGAGACCCGGTATAGCCCCCATCCCTCCC
 310 320
 SerHisHisGlnLeuThrLeuProAspProSerHisHisGlyLeuHisSerThrProAsp
 AGCCACCACTGACTCTCCCCGACCCGTCCTCCACCATGGTCTCCACAGCACTCCCGAC
 IIIIIIIIIII IIIIIIIII IIIIIIIII IIIIIIIII IIIIIIIII IIIIIIIII IIIIIIIII IIIIIIIII
 AGTCACCACTGACTCTCCCTGATCCATCCACCGGCTACATAGCAACCTGAC
 Asn 960
 330 339
 SerProAlaLysProGluLysAsnGlyHisAlaLys AspHisProLysIleAlaLys
 AGCCCCGCCAAACAGAGAAGAATGGGCATGCCAAA...GACCACCCCAAGATTGCCAAG
 IIIIIIIIIII IIIIIIIIIII IIIIIIIII IIIIIIIII IIIIIIIII IIIIIIIII IIIIIIIII IIIIIIIII
 AGTCCTGCCAAACAGAGAAGAATGGGCACGCCCAAGATTGTCAATCCCAGGATTGCCAAG
 IleValAsn Arg

FIG. 18-6

SUBSTITUTE SHEET

IlePheGluIleGlnThrMETProAsnGlyLysThrArgThrSerLeuLysThrMETSer
 ATCTTTGAGATCCAGACCATGCCCAATGGCAAACCCGGACCTCCCTCAAGACCATGAGC
 TTTCTTGAGATCCAGACCATGCCCAATGGCAAACCCGGACCTCCCTTAAAGACGATGAGC
 phe

ArgArgLysLeuSerGlnGlnLysGluLysLysAlaThrGlnMETLeuAlaIleValLeu
369 379

[illegible]

FIG. 18-7

41 / 47

CAGGCCGGCCAGCCTCACCCCTTGCGAACCGTGAGCAGGAAGGCCCTGGGTGGATCGGCCCTC 1437

CTCTTCTTAGCCCCGGCAGGCCCTGCAGTGTTCGCTTGCTCCATGCTCCTCACTGCCCG

CACACCCCTCACTCTGCCAGGGCAGTGCTAGTGAGCTGGGCATGTTACCAAGCCCTGGGGCT 1557

GGCCCCAGCTCAGGGGCAGCTCATAGAGTCCCCCTCCACCTCCAGTCCCCCTATCCTT

GGACCAAGATGCAGCGCCCTTCCTTGACCTTCCTCTGGGGCTCTAGGGTTGCTGGAGC 1677

CTGAGTCAGGGCCAGAGGCTGAGTTTCTCTTTGTGGGGCTTGGCGTGGAGCAGCGCGT

GGGGAGAGATGGACAGTTACACCCCTGCAAGGCCCCACAGGAGGCAAGCAAGCTCTCTTGC 1797

CGAGGAGCCAGGCAACTTCAGTCTCTGGGAGACCCATGTAAATACCAGACTGCAGGTTGGA

CCCCAAGGATTCCCAGCCAAACCTTAGCTCCCTCCCGCACCCCGATGTGGACCTCTA 1917

FIG. 18-9


SUBSTITUTE SHEET

42 / 47

CTTCCAGGCTAGTCCGGACCCACCTCACCCCGTTACAGCTCCCAAGTGTTCCACAT
GCTCTGAGAAGAGAGCCCTCATCTTGAAGGCCCCAGGAGGTCTATGGGAGAGGAACT 2037
CCTTGGCCTAGCCACCCCTGCTGCCTTCTGACGGCCCTGCAATGTATCCCTTCTCACAGC
ACATGCTGGCCAGCCTGGGGCCTGGCAGGGAGGTCAGGCCCTGGAACCTATCTGCGCCT 2157
GGGCTAGGACATCAGAGGTTCTTTGAGGGACTGCCCTCTGCCACACTCTGACGCAAACC
ACTTTCCTTTTCTATTCCCTTCTGGCCCTTTCCTCTCTCCTGTTTCCCTTCCCTTCCACTGC 2277
CTCTGCCTTAGAGGAGCCACGGCTAAGAGGCTGCTGAAAACCATCTGGCCTGGCCTGGC
CCTGCCCTGAGGAAGAGGGCAAGCTGCAGCTTGGGAGAGCCCCTGGGGCCTAGACTCTG 2397
TAACATCACTATCCGATGCACCAAACTAATAAACTTTGACGAGTCACCTTC (A)_n 2449

FIG. 18-10**SUBSTITUTE SHEET**

43 / 47



SUBSTITUTE SHEET

44 / 47

1 2 3 4

kb

—12.0

— 6.0

— 4.0

— 2.0


— 1.6

— 1.0

FIG. 20

45 / 47

FIG. 21



46 / 47

DRUG	HUMAN D ₂	RAT D ₂	RAT STRIATUM
PIPERONE	0.125	0.35	0.56
(+) BUTACLAMOL	0.94	1.2	1.6
HALOPERIDOL	2.4	5.1	5.8
SULPIRIDE	206	160	205
MIANSERIN (5-HT)	2685	4300	4600
SCH 23390 (DI)	2145	2500	3300
(+) BUTACLAMOL	>10,000	>10,000	>10,000
Kd [³ H] DOMPERIDONE	0.74	0.40	0.40

FIG. 22

47 / 47

IG. 23

285 ¹	286
exon1...GGAGgtagtg...intron1...tccccagGTGG...exon2	
395	396
exon2...ACAGgtgagcc...intron2...cttgccagGTAC...exon3	
532	533
exon3...GCAGgtacatt...intron3...ccccccagACCA...exon4	
723	724
exon4...AAAGgtctctcaa...intron4...tccacagGGCA...exon5	
810	811
exon5...AGTGgtaagtg...intron5...gggtgcagGAGG...exon6	
1138	1139
exon6...CTCGgtgagtc...intron6...ccccccagGCGT...exon7	

¹Numbering begins with A of the putative initiator methionine codon (see Fig. 18)

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US89/05285

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC IPC(5): C12N 15/00 C12N 7/01 C12N 5/10 C12P 21/02 U.S. Cl.: 536/27 435/235 435/240.1 435/68		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
U.S.	536/27 435/235, 320, 240.1, 68	
Documentation Searched other than Minimum Documentation: to the Extent that such Documents are Included in the Fields Searched ⁸		
Genbank EMBL and PIR databases searched for sequence in figure 1. Dialog database Keywords searched: mammal? D2 dopamine receptor DNA probe expression		
III. DOCUMENTS CONSIDERED TO BE RELEVANT ⁹		
Category [*]	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
Y	US, A, 4,705,858 HADLEY 10 November 1987. See column 19 lines 49-54.	3-4, 21-22
Y	Nature, Volume 329, Issued 3 September 1987 KOBILKA "An intronless gene encoding a potential member of the family of receptors coupled to guanine nucleotide regulating proteins" See figures 1 and 2 (IVFGNVLV peptide). and pages 75-79	3-4, 21-22
X Y	Nature, Volume 321, Issued 01 May 1986 Dixon, "Cloning of the gene and cDNA for mammalian Beta-adrenergic receptor and homology with rhodopsin" see pages 75-79. See figure 4 peptide SSIVSFYVP and figure 3 legend regarding labelled oligonucleotide.	1-2,7-8 11-17, 19-20 & 53 3-6,9-10, 18,21-22
<p>[*] Special categories of cited documents: ¹⁰</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"A" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
14 February 1990	28 FEB 1990	
International Searching Authority	Signature of Authorized Officer	
ISA/US	MICHELLE S. MARKS-PATENT EXAMINER	

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)

Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No
<u>X</u> <u>Y</u>	<u>The Journal of Biological Chemistry</u> , Volume 262(15) Issued 25 May 1987 KOBILKA "Delineation of the intronless nature of the genes of the human and hamster Beta 2 - Adrenergic Receptor and their putative promoter regions". See figures 2, 5 and 9 and pages 7321-7327.	1-2,5-9; <u>53</u> 3-4,11-22
<u>X,P</u> <u>Y,P</u>	<u>Nature</u> , Volume 336 Issued 22/29 December 1988 BUNZOW "Cloning and expression of a rat DZ dopamine receptor cDNA" see pages 783-787 and figure 1.	1-2,5-9 <u>53</u> 3-4,11-22

FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

V. ☐ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE¹

This international search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. ☐ Claim numbers _____, because they relate to subject matter¹² not required to be searched by this Authority, namely:

2. ☐ Claim numbers _____, because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out¹³, specifically:

3. ☐ Claim numbers _____, because they are dependent claims not drafted in accordance with the second and third sentences of PCT Rule 6.4(a).

VI. ☒ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING²

This International Searching Authority found multiple inventions in this international application as follows:

See Attachment.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.

2. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:

3. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers: 1-9, 11-22 and 53

as per telephonic Restriction Practice

4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the International Searching Authority did not invite payment of any additional fee.

Remark on Protest

☐ The additional search fees were accompanied by applicant's protest.

☐ No protest accompanied the payment of additional search fees.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.